

Short communication

First report of *Anarsia (Ananarsia) eleagnella* (Lep.: Gelechiidae: Dichomeridinae) from Iran

G. A. Asadi^{1&*}, U. Schaffner², M. Cristofaro³, V. Ghasemi⁴, S. Y. Sinev⁵ and R. Ghorbani¹

1. Department of Agronomy, Faculty of Agriculture, Ferdowsi University of Mashhad, P.O. Box 91775-1163, Mashhad, Iran, 2. CABI Europe-Switzerland, Rue des Grillons 1, 2800 Delémont, Switzerland, 3. BBICA, Via del Bosco 10, 00060 Sacrofano (Rome), Italy, 4. Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran, 5. Russian Academy of Sciences, Zoological Institute RAS, Saint Petersburg, Russia.

*Corresponding author, E-mail: asadi@um.ac.ir

چکیده

در بررسی عوامل کنترل‌کننده طبیعی علف‌های هرز ایران که طی سال‌های ۱۳۸۹ تا ۱۳۹۱ صورت گرفت، میوه‌های سنجد حاوی لارو یک گونه شب‌پره، در شهرستان‌های واقع در استان‌های خراسان شمالی، خراسان رضوی، خراسان جنوبی، آذربایجان شرقی، آذربایجان غربی و کردستان مشاهده و جمع‌آوری گردید. پس از پرورش لاروها در آزمایشگاه، حشرات کامل با نام علمی *Anarsia (Ananarsia) eleagnella* Kuznetsov شناسایی شدند که توسط نگارنده پنجم نیز مورد تأیید قرار گرفت. این گونه برای اولین بار از ایران گزارش می‌شود.

The plant species *Elaeagnus angustifolia* L. is native to western and central Asian countries including south of Russia, Kazakhstan as well as Turkey and Iran. It is considered as an invasive species in Europe and America (Christensen, 1963; Katz & Shafroth, 2003). *Anarsia eleagnella* Kuznetsov (fig. 1, A) is a widely distributed lepidopteran species being recorded from Romania, southern parts of Ukraine and European Russia, Transcaucasia, Turkmenistan, Kazakhstan, Afghanistan, Hungary and southern Siberia (Szabóky *et al.*, 2009). In addition to *E. angustifolia*, the plant species of the genus *Elaeagnus* and probably *Hippophae* of the family Elaeagnaceae are the hosts of *A. eleagnella* (Ponomarenko, 1997, 1999).

In a survey conducted on the identification of the biological control agents of weeds in Iran, from 2010 to 2012, the infested fruits of *E. angustifolia* containing moth larvae were collected from the provinces of Khorasan-e-Shomali, Khorasan-e-Razavi, Khorasan-e-Jonubi, Azarbaijan-e-Sharghi, Azarbaijan-e-Gharbi and Kordestan. The larvae (fig. 1, B) were reared in growth chamber until adult moths emerged. The attacking moth was identified as *Anarsia (Ananarsia) eleagnella* Kuznetsov, 1957 which was later confirmed by the fifth author. This species is a new record for the insect fauna of Iran. Previous to this record only a single pest species of the genus *Anarsia* Zeller, peach twig borer (*Anarsia lineatella* Zeller, 1839), had been reported

from Iran (Esmaili, 1997). The species *A. eleagnella* belongs to the tribe Ananarsiini and subfamily Dichomeridinae. The specimens are deposited in the insect collection of the Department of Agronomy, Faculty of Agriculture, Ferdowsi University of Mashhad, Iran.

The adult moth is grayish brown, about 4.5 mm long and an average wingspan of 11 mm (n = 7), marked with numerous brown spots on the forewing, especially at apex; hindwings with less markings, fringed with relatively long hairs in both front and hind margins; antenna gray, setiform, almost half of body length (fig. 1, A). Pupa 2.5-3.5 mm, brown, without a cocoon (n = 6) (fig. 1, F).

Preliminary biological studies indicate that *A. eleagnella* has two generations in Mashhad (Asadi *et al.*, unpublished data) where the first generation larvae feed on terminal shoots (fig. 1, C) and the second generation attack unripe fruit (fig. 1, D-E) causing heavily infested fruits fall off trees.




Fig. 1. *Anarsia eleagnella*: (A) adult, (B) larvae, (C) shoot strike damage caused by overwintered larvae, (D-F) damaged fruit caused by second generation larvae, (E) pupae.

References

- Christensen, E. M.** (1963) Naturalization of Russian olive (*Elaeagnus angustifolia* L.) in Utah. *The American Midland Naturalist* 70(1), 133-137.
- Esmaili, M.** (1997) *Fruit trees pests*. 578 pp. Sepehr Publication Center, Tehran.
- Katz, G. L. & Shafroth, P. B.** (2003) Biology, ecology, and management of *Elaeagnus angustifolia* L. (Russian olive) in western North America. *Wetlands* 23(4), 763-777.
- Ponomarenko, M. G.** (1997) Catalogue of the subfamily Dichomeridinae (Lepidoptera, Gelechiidae) of the Asia. *Far Eastern Entomologist* 50, 1-67.
- Ponomarenko, M.G.** (1999) Subfamily Dichomeridinae. pp. 194-257 in Lerh, P.A. (Ed.) *Key to the insects of the Russian Far East; Vol. 5, Part 2: Lepidoptera and Trichoptera*. 672 pp. Dalnauka Publishing.
- Szabóky, C. S., Tokár, Z., Liška, J. & Pastorális, G.** (2009) New data to the Microlepidoptera fauna of Hungary, part XII (Lepidoptera: Lypusidae, Bucculatricidae, Yponimeutidae, Depressariidae, Coleophoridae, Blastobasidae, Autostichidae, Gelechiidae, Tortricidae). *Folia Entomologica Hungarica* 70, 139-146.

Received: 5 January 2013

Accepted: 24 February 2013