

Short communication

First record of *Alloxysta pleuralis* from Iran (Hym.: Cynipoidea: Figitidae: Charipinae)

M. Ferrer-Suay<sup>1&\*</sup>, J. Selfa<sup>2</sup>, Z. Feli Kohikheili<sup>3</sup>, M. R. Damavandian<sup>3</sup> and J. Pujade-Villar<sup>1</sup>

1. Universitat de Barcelona, Facultat de Biologia, Departament de Biologia Animal. Avda. Diagonal 645, 08028-Barcelona, Spain, 2. Universitat de València, Facultat de Ciències Biològiques, Departament de Zoologia, Campus de Burjassot-Paterna, Dr. Moliner 50, E-46100 Burjassot (València), Spain, 3. Department of Plant Protection, Sari Agricultural Sciences and Natural Resources University, P.O. Box 578, Sari, Iran.

\*Corresponding author, E-mail: mar.ferrer.suay@gmail.com

چکیده

طی نمونه برداری های انجام شده از زنبورهای پارازیتوبید در استان مازندران، گونه *Alloxysta pleuralis* (Cameron, 1879) از *Citrus aurantium* L. (Rutaceae) به عنوان پارازیتوبید شته *Aphis gossypii* Glover درخت نارنج، در شهرستان جویبار جمع آوری و شناسایی شد. این زنبور پارازیتوبید برای اولین بار از ایران گزارش می شود.

As the result of a field study based on the parasiting wasps of important citrus pests in Mazandaran province, Iran; a species of the genus *Alloxysta* Förster (Hym.: Cynipoidea: Figitidae: Charipinae) was collected. This genus is one of the eight genera that compose the subfamily Charipinae. The other seven genera are as follows: *Apocharips* Ferguson, *Dilapothor* Paretas-Martínez & Pujade-Villar, *Dilyta* Förster, *Lobopterocharips* Paretas-Martínez & Pujade-Villar, *Lytoxysta* Kieffer, *Phaenoglyphis* Förster and *Thoreauana* Girault. Members of the Charipinae are very small with smooth and shiny bodies, and small interspecific variability. They are widely distributed around the world.

*Alloxysta* is the most diverse genus within Charipinae, including 111 species (Ferrer-Suay *et al.*, 2012). Species of *Alloxysta* are hyperparasitoids of aphids (Hemiptera: Aphididae) via Aphidiinae (Hym.: Ichneumonoidea: Braconidae) and Aphelininae (Hym.: Chalcidoidea: Aphelinidae).

The study of Iranian Charipinae has been significantly improved in the recent years. Ferrer-Suay *et al.* (2013b) identified 15 species, of which the following nine species were first recorded from Iran: *Alloxysta arcuata* (Kieffer, 1902), *A. brevis* (Thomson, 1862), *A. castanea* (Hartig, 1841), *A. macrophadna* (Hartig, 1841), *A. melanogaster* (Hartig, 1840), *A. pusilla* (Kieffer, 1902), *A. ruficollis* (Cameron, 1883), *A. tscheki* (Giraud, 1860) and *A. ullrichi* (Giraud, 1860). Later, the species *A. ramulifera* (Thomson,

1862) was added to the list of charipine fauna of Iran (Ferrer-Suay *et al.*, 2013a).


In order to collect and identify the parasitoid wasps of Caspian coast province of Mazandaran, the aphid infested branches were collected from 11 towns and transferred to the laboratory for rearing parasitoids during 2011-13. The emerged parasitoid wasps were transferred to 75% ethanol upon their emergence. Later, the species *Alloxysta pleuralis* (Cameron, 1879) was identified as a parasitoid of *Aphis gossypii* Glover, (Hem.: Aphididae) on *Citrus aurantium* L. (Rutaceae). This is the first report of *A. pleuralis* from Iran.

A total of nine *Alloxysta* specimens were collected in two samples of *C. aurantium* from Juybar (Mazandaran) on 23.4.2013. The specimens were superficially very close to those of *A. pleuralis*. This species is easily distinguished from other *Alloxysta* species by the following combination of characters: partially open radial cell (fig. 1, a); pronotal carinae present; two propodeal carinae reaching the base independently (fig. 1, b); female antenna: F1 longer than F2, F2 shorter than F3 and F3 shorter than F4 (fig. 1, c-d); male antenna: F1-F3 subequal in length and slightly curved.

The specimens have been deposited at the University of Barcelona: "Ex. *Citrus* sp., with Coccoidea, col. 23.4.2013", "Mazandaran (Iran), Leg. Zahra Feli", "*Alloxysta pleuralis* (Cameron, 1879) ♀ M. Ferrer-Suay det. 2013": (2 ♀♀).

*Alloxysta pleuralis* is widely distributed in the Palaearctic region and has been reported from England, France, Germany, Ireland, Israel, Norway, Poland,

Scotland and Spain (Ferrer-Suay *et al.*, 2012). It has also been recorded from India (Ahmad & Singh, 1996).



**Fig. 1.** *Alloxysta pleuralis*: (a) forewing and detail of the radial cell, (b) propodeal plate, (c) detail of the female antenna, (d) female antenna.

## References

- Ahmad, Md. E. & Singh, R. (1996) Records of aphid parasitoids from the Bihar and associations with their hosts and food plants. *Journal of Advanced Zoology* 17(1), 26-33.
- Cameron, P. (1879) On some new or little known British Hymenoptera. *Transactions of the Entomological Society of London* 1879, 107-119.
- Cameron, P. (1883) Descriptions of sixteen new species of parasitic Cynipidae, chiefly from Scotland. *Transactions of the Entomological Society of London* 16(4), 365-374.
- Ferrer-Suay, M., Paretas-Martínez, J., Selfa, J., & Pujade-Villar, J. (2012) Taxonomic and synonymic world catalogue of the Charipinae and notes about this subfamily (Hymenoptera: Cynipoidea: Figitidae). *Zootaxa* 3376, 1-92.
- Ferrer-Suay, M., Selfa, J., Safoora, F., Karimi, J. & Pujade-Villar, J. (2013a) First records of *Alloxysta ramulifera* (Thomson, 1862) and *Asaphes vulgaris* Walker, 1834 from Iran. *Linzer biologische Beiträge* 45(1), 671-672.
- Ferrer-Suay, M., Selfa, J., Seco-Fernández, M. V., Melika, G., Alipour, A., Rakhshani, E., Talebi, A. A. & Pujade-Villar, J. (2013b) A contribution to the knowledge of Charipinae (Hymenoptera: Cynipoidea: Figitidae) associated with aphids from Iran, including new records. *North-Western Journal of Zoology* 9(1), 30-44.
- Giraud, J. (1860) Enumeration des Figitides de l'Autriche. *Verhandlungen der kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien* 10, 123-176.
- Hartig, T. (1840) Ueber die Familie der Gallwespen. *Zeitschrift für Entomologie (Germar)* 2, 176-210.
- Hartig, T. (1841) Erster nachtrag zur naturgeschichte der Gallwespen. *Zeitschrift für Entomologie (Germar)* 3, 322-358.
- Kieffer, J. J. (1902) Les Cynipides (part 2). In: Andre, E. *Species des Hyménoptères d'Europe et d'Algérie*, 7(2): 748 pp. + 21 pl. [Charipinae in: 5-78 + 592-602 (= 1904a)].
- Thomson, C. G. (1862) Forsok till uppställning och beskrifning af Sveriges Figiter. *Öfversigt af Kongl. Svenska Vetenskaps-akademiens förhandlingar* 18, 395-420.

Received: 21 February 2014

Accepted: 19 September 2014