





Short Communication

The occurrence of the citrus flower moth, *Prays citri* (Lepidoptera: Yponomeutoidea: Praydidae) for the first time in Iran

Helen Alipanah¹ , Mohammad Reza Damavandian² , Maedeh Abdi³  & Lida Fekrat³ 

1- Insect Taxonomy Research Department, Iranian Research Institute of Plant Protection (IRIPP), Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

2- Department of Plant Protection, Faculty of Crop Sciences, Sari Agricultural Sciences and Natural Resources University, Sari, Iran

3- Department of Plant Protection, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

Abstract. The citrus flower moth, *Prays citri* (Millière), was collected in Chubāgh village, Sari, Mazandaran Province, Iran, using ferolite water-based traps on Lemon trees at a greenhouse in May and July of 2024. The considerable damages on Lemon flowers caused by this pest were observed at the same time. This is the first report of the genus and species from Iran, which was previously considered as an external quarantine pest in the country. A brief taxonomic characterization of the species, as well as the adult and genitalia figures of the examined specimens, are provided.

Article history

Received: 03 September 2024

Accepted: 08 October 2024

Published: 08 January 2025

Subject Editor: Ebrahim Ebrahimi

Corresponding author: Helen Alipanah

E-mail: halipanah@gmail.com

DOI: <https://doi.org/10.61186/jesi.45.1.12>

Keywords: Citrus, external quarantine pest, Lemon, new record, Iran

The genus *Prays* Hübner, 1825, the largest genus of the family Praydidae, consists of 59 described species worldwide (Cong & Li, 2017; Bippus, 2020), with the most diversity in Asia (Cong & Li, 2017). This genus is characterized by the absence of pectens on the scape of the antennae, the membranous eighth tergum of the male not protruded medially on the posterior margin, the presence of the sacculus process in male genitalia, and a rudimentary uncus (Sohn & Wu, 2011; Cong & Li, 2017).

In May and July of 2024, several moth specimens were collected using ferolite water-based traps on Citrus trees at a greenhouse in Sari, Mazandaran Province, and identified as *Prays citri* (Millière, 1873) following morphological and genitalia examination. The considerable damages on Lemon flowers caused by their larvae were observed concurrently. This species, which was previously being considered an external quarantine pest in Iran, has historically been a sporadic and minor pest on the flowers of Lemons and Limes. However, in recent years, it has become a regular pest, particularly causing damage to fruit up to a golf-ball size (Grout & Moore, 2024). The genus and species here are being reported from Iran for the first time.

Prays citri (Millière, 1873)

Material examined

Māzandarān Prov.: 3 ♂♂ 5 ♀♀, Sāri, Juybār County, Chubāgh village (Mr. Shaeri greenhouse), 36°45'21"N, 52°58'27"E, -23 m, 12.v.2024, 3.vii.2024, Damavandian legs (genitalia slides HA-2978, HA-2979, HA-2980, HA-2981, HMIM; HA: Helen Alipanah; HMIM: Hayk Mirzayans Insect Museum).

Morphological characters

Male, female similar externally, female somewhat lighter. Forewing length 5.2–6.2 mm; wingspan 11.5–14.6 mm. Forewing greyish-brown, strongly mottled by spots, costal margin with few ill-defined blackish-brown strigulae alternated with whitish-grey on basal half; fringes greyish-brown. Hindwing brownish-grey, fringes concolorous (Fig. 1E) (Tamutis *et al.*, 2022; Grout & Moore, 2024). There was no external variation among the studied material.




Fig.1. Last larval instar and its damage (A), damages caused by the larvae (B, C), pupa inside cocoon (D), and the adult male of *Prays citri* (Millière), upperside. Arrows in the figures B and C indicate the damages caused by the larvae on sepals of the Citrus flower in the greenhouse in Sari region, Mazandaran, Iran. (A, after CABI, 2024; B–E, original; photos: B–D, Mohammad Reza Damavandian, E, Helen Alipanah).

Male genitalia (Fig. 2A–G) with uncus as a small swelling (Fig. 2C, F), sometimes hardly visible (Fig. 2A); socius moderate to relatively short (based on slide preparation) (Fig. 2A, C, F), bearing long setae (Fig. 2A); tegumen trapezoidal; gnathos stripe-shaped, narrow, with a small, nearly triangular projection posteriorly; valva in the examined specimens finger-shaped at the end of cucullus, sacculus broadened towards the base, with a tuft of strong dense setae and a relatively long, acute thorn distally; anellus lobes with long setae; vinculum V-shaped, broad; saccus rounded apically; phallus slender, curved medially (Tamutis *et al.*, 2022), in the examined specimens 1.66–1.67 times as long as the valva ($n = 2$), vesica with two long rod-shaped cornuti (Fig. 2A, B) and rows of minute spines. In another specimen, there seems to be three rod-shaped cornute in vesica (Fig. 2C, D).

Female genitalia (Fig. 2H–L) with nearly triangular papillae anales; apophyses posteriores slightly longer than apophyses anteriores; lamella postvaginalis produced into a pair of finger-like sclerotized plates, with sparse long setae posteriorly; sternum VIII with longitudinal central groove; antrum short, cup-shaped; ductus bursae relatively short with sclerotized wall; corpus bursae oviform, signum placed medially, elliptical, concave elliptically at posterior end, with dentate margins (Tamutis *et al.*, 2022).

Distribution

Prays citri is widespread in the Mediterranean region. It is present in Africa, Asia, Oceania, and Europe (CABI, 2024). According to CABI (2024), the identifications of *P. citri* on Citrus from east of Turkey and the Middle East (Sri Lanka, Malaysia, Philippines, Pakistan, Fiji, Samoa) are likely to be erroneous.

Biology

The larvae (Fig. 1A) vary from light grey or green to brown with a brown head (Grout & Moore, 2024). The chrysalis (Fig. 1D) is found in a very soft, white and sharp cocoon (Bissanti, 2023), and the pupa is initially light green but soon turns chocolate brown (Grout & Moore, 2024). The adults fly at night and twilight. The females lay 1-3 eggs in spring preferably on the petals of the still closed flowers, and also in the sepals, shoots, or small fruits.

The last larval instars normally pupate inside the flowers in which they have already fed (Bissanti, 2023). They overwinter as chrysalis, but sometimes in the pupal and larval stages, or as eggs or adults (Bissanti, 2023; Tamutis *et al.*, 2022). They can complete at least three (Karamaouna *et al.*, 2009), and up to 11, annual generations (Badr *et al.*, 2018).




Fig. 2. Male (A–G) and female (H–L) genitalia of *Prays citri* (Millière): A, C) Main body and phallus in ventral and lateral views, respectively (C, with pleural lobes of eighth abdominal segment). B, D) Distal one-third of the phallus. E) Pleural lobes of eighth abdominal segment. F) Uncus and socii. G) Distal end of the valva and sacculus. H, K) Main body in ventral view. I, J) Signum. L) Lamella postvaginalis and sternum VIII (genitalia slides, HA-2978, HA-2979, HA-2980, HA-2981, HMIM; slide preparation and photos: Helen Alipanah).

Host plants

Citrus species (Rutaceae) are the main host plants of *Prays citri*. However, it prefers Lemon (*Citrus limon* (L.)), Key lime (*C. ×aurantifolia* Swingle), Pummelo (*C. decumana* L.), Mandarin orange / tangerine (*C. reticulata* Blanco), Sweet orange (*C. ×sinensis* (Osbeck)), and Grapefruit (*C. ×paradisi* Macfad.) (Ibrahim & Shahateh, 1984; Abd El-Kareim et al., 2017; Tamutis et al., 2022). The other known host plants of the species are White sapote, Sour orange, *Citrus microcarpa* Bunge (Rutaceae), Broad-leaf privet (Oleaceae), and Sapodilla (Sapotaceae) (Sinacori & Mineo, 1997; Martinez et al., 2019; Tamutis et al., 2022; CABI, 2024).

Damage

The larvae are particularly destructive to blossoms and young fruit which occur in spring and mid-summer. Damaged flower buds and flowers quickly wilt and are shed (Grout & Moore, 2024). Penetration marks of the larvae are initially fairly small, but once the fruit matures, they appear as brown necrotic scars (Moore & Kirkman, 2014) (Fig. 1B, C). In the center of these spots, the chorion of the egg can be found.

Author's Contributions

Helen Alipanah: conceptualization, investigation, draft preparation, slide preparation and photography of the adults and genitalia, final review and edit. **Mohammad Reza Damavandian:** investigation, collecting of specimens, photography of the symptoms and cocoon, final review and edit. **Maedeh Abdi:** investigation and collecting of specimens. **Lida Fekrat:** final review and edit.

Author's Information

Helen Alipanah	✉ halipanah@gmail.com	 https://orcid.org/0000-0002-3717-6304
Mohammad Reza Damavandian	✉ damavandianm@gmail.com	 http://orcid.org/0000-0002-4403-569X
Maedeh Abdi	✉ abdi.maedeh@mail.um.ac.ir	 https://orcid.org/0009-0003-6886-1739
Lida Fekrat	✉ fekrat@um.ac.ir	 https://orcid.org/000-0002-5189-9488

Funding

This work is supported financially by the Iranian Research Institute of Plant Protection (IRIPP), Ferdowsi University, Mashhad, Agriculture Faculty, and Sari Agricultural Science and Natural Resources University.

Data Availability Statement

The specimens examined in this study are deposited in the Hay Mirzayans Insect Museum (HMIM) and are available by the curator upon request.

Acknowledgments

We express our thanks to Eng. Mohammad Reza Nematian (Iranian Research Institute of Plant Protection (IRIPP), Tehran, Iran) for spreading the specimens. Dr Milad Habibi and Eng. Mohammad Reza Jafari (Sari Agricultural Sciences and Natural Resources University, Sari, Iran) are acknowledged for their attempts in sending the specimens to IRIPP, Tehran.

Ethics Approval

Insects were used in this study. All applicable international, national, and institutional guidelines for the care and use of animals were followed. This article does not contain any studies with human participants performed by the author.

Conflict of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

REFERENCES

- Abd El-Kareim, A. I., Moustafa, S. A., Fathy, D. M. & Amara, T. M. (2017) Ecological and biological studies on the citrus flower moth, *Prays citri* (Millière) (Lepidoptera: Hyponomeutidae). *Journal of Plant Protection and Pathology* 8 (8), 397–402. <https://doi.org/10.21608/jppp.2017.46353>
- Badr, A. F., Hekal, A. M. & Youssef, L. A. (2018) Population fluctuations of immature stages of the citrus flower moth, *Prays citri* Mill. (Lep.: Hyponomeutidae) on lime trees. *Arab Universities Journal of Agricultural Sciences* 26 (spec. iss. 2C), 1993–2000. <https://doi.org/10.21608/ajs.2018.31666>
- Bippus, M. (2020) Records of Lepidoptera from the Malagasy region with description of new species (Lepidoptera: Tortricidae, Noctuidae, Alucitidae, Choreutidae, Euteliidae, Gelechiidae, Blastobasidae, Pterophoridae, Tonzidae, Tineidae, Praydidae, Cosmopterigidae, Batrachedridae). *Phelsuma* 28, 60–100.
- Bissanti, G. (2023) Un Mondo Ecosostenibile, dentro i codici della Natura. *Prays citri*. Available from: <https://antropocene.it/en/2023/02/08/prays-citri-2/> (accessed 8 August 2024).
- CABI (2024). *Prays citri* (citrus flower moth). Invasive species Compendium. Wallingford, UK. Available from: <https://www.cabi.org/isc/datasheet/43910> (accessed 12 August 2024).
- Cong, P. & Li, H. (2017) Review of the genus *Prays* Hübner, 1825 (Lepidoptera: Praydidae) from China, with descriptions of twelve new species. *Zootaxa* 4263 (2), 201–227. <https://doi.org/10.11646/zootaxa.4263.2.1>

- Grout, T. G. & Moore, S. D. (2024) Flower moth (Citrus). Text extracted with permission from the editors from: Prinsloo, G. L. & Uys, V. M. (Eds) 2015. *Insects of Cultivated Plants and Natural Pastures in Southern Africa*. Entomological Society of Southern Africa. Available from: <https://insectscience.co.za/pest/citrus-flower-moth> (accessed 10 August 2024).
- Ibrahim, S. S. & Shahateh, W. A. (1984) Biological studies on the citrus flower moth *Prays citri* Miller. (Lep., Hyponomeutidae) in Egypt. *Arab Journal of Plant Protection* 2 (1), 4–9 [in Arabic].
- Karamaouna, F., Mylonas, P., Papachristos, D., Kontodimas, D., Michaelakis, A. & Kapaxidi, E. (2009) Main Arthropods pest of Citrus culture and pest management in Greece. Ciancio, A. & Mukerji, K. G. (Eds), *Integrated Management of Arthropod Pests and Insect Borne Diseases*, Integrated Management of Plant Pests and Diseases; Springer: Dordrecht, The Netherlands, 29–59. https://doi.org/10.1007/978-90-481-8606-8_2
- Martinez, J. I., Crane, J. H., Wasielewski, J., Miller, J. Y. & Carrillo, D. (2019) Lepidoptera pest of sapodilla (*Manilkara zapota* (L.) van Royen) in south Florida, with some comments on life history and natural control. *Insecta Mundi* 739 (739), 1–26. <https://doi.org/10.5281/zenodo.3676599>
- Moore, S. D. & Kirkman, W. (2014) The Lemon Borer Moth = The Citrus Flower Moth, *Prays citri*: Its biology and control on citrus. *The South African fruit Journal / Die Joernaal vir die Vrugtebedryf in Suid-Afrika* 13 (1), 86–91.
- Sinacori, A. & Mineo, N. (1997) Two new host plants of *Prays citri* and *Contarinia* sp. (?) *citri* [*Casimiroa edulis* – *Ligustrum lucidum* – Sicily]. *Informatore Fitopatologico* 47 (7–8), 13–15 [in Italian].
- Sohn, J.-C. & Wu, C.-Sh. (2011) A taxonomic review of *Prays* Hübner, 1825 (Lepidoptera, Yponomeutoidea, Praydidae) from China with description of two new species. *Tijdschrift voor Entomologie* 154, 25–32. <https://doi.org/10.1163/22119434-900000306>
- Tamutis, V., Sruoga, V., Česonienė, L. & Daubaras, R. (2022) The first appearance of the citrus flower moth (*Prays citri* (Millière, 1873)) (Lepidoptera: Praydidae) in Lithuania. *Zemdirbyste-Agriculture* 109 (1), 89–93. <https://doi.org/10.13080/z-a.2022.109.0012>





Citation: Alipannah, H., Damavandian, M. R., Abdi, M. & Fekrat, L. (2025) The occurrence of the citrus flower moth, *Prays citri* (Lepidoptera: Yponomeutoidea: Praydidae) for the first time in Iran. *J. Entomol. Soc. Iran*, 45 (1), 145–150.

DOI: <https://doi.org/10.61186/jesi.45.1.12>

URL: https://jesi.areeo.ac.ir/article_131275.html



گزارشی از وجود شب‌پره گل مرکبات، *Prays citri* (Lepidoptera: Yponomeutoidea: Praydidae) برای اولین بار در ایران

هلن عالی پناه^۱ , محمدرضا دماندیان^۲ , مانده عبدی^۳  و لیدا فکرت^۳ 

- ۱- بخش تحقیقات رده‌بندی حشرات، مؤسسه تحقیقات گیاه پزشکی کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران
- ۲- گروه گیاهپزشکی، دانشکده علوم زراعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ایران
- ۳- گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه فردوسی مشهد، مشهد، ایران

چکیده: در اردیبهشت و تیر ماه جاری نمونه‌هایی از شب‌پره گل مرکبات، *Prays citri* (Millière) در تله‌های فرولیت آبی کار گذاشته شده در گلخانه‌ای در روستای چوبگاه شهرستان ساری، استان مازندران، ایران، جمع‌آوری شدند و در همان زمان، خسارت قابل توجه ایجاد شده توسط لاروهای این آفت روی گل‌های لیمو مشاهده شد. جنس و گونه این آفت که تا به حال به‌عنوان آفت قرنطینه خارجی در کشور در نظر گرفته می‌شد، برای اولین بار از ایران گزارش می‌شوند. در این مقاله مشخصات تاکسونومیک گونه به‌همراه شکل حشره کامل، ساختار اندام‌های تناسلی نر و ماده نمونه‌های مورد بررسی، و نحوه خسارت آن‌ها به اختصار ارائه شده‌اند.

تاریخچه مقاله
دریافت: ۱۴۰۳/۰۶/۱۳
پذیرش: ۱۴۰۳/۰۷/۱۷
انتشار: ۱۴۰۳/۱۰/۱۹
دبیر تخصصی: ابراهیم ابراهیمی
نویسنده مسئول: هلن عالی پناه

پست الکترونیک: halipanah@gmail.com

DOI: <https://doi.org/10.61186/jesi.45.1.15>

کلمات کلیدی: آفت قرنطینه خارجی، گزارش جدید، ایران، Lemon، Citrus