

Research Article

Checklist of beetles (Coleoptera) from the cork oak forest of Larache (Morocco)

Amine Samih^{1,2} , Nouredin Maatouf² , Sergi Trócoli³ , Hamza Habbaz^{1,2}  & Latifa Rohi¹ *1- Laboratory of Ecology and Environment, Faculty of Sciences Ben M'sik, University of Hassan II, Casablanca, Morocco**2- Center for Innovation, Research, and Training, National Agency for Water and Forests, Rabat, Morocco**3- Museu de Ciències Naturals de Barcelona, Laboratori de Natura, Col·lecció d'artròpodes, Barcelona, Spain*

Abstract. The cork oak forest of Larache, a vital part of Morocco Atlantic coastal cork oak woodlands, plays a crucial role in supporting local ecosystems and human societies. However, it is increasingly threatened by intensive agriculture and other anthropogenic pressures, resulting in habitat loss for insects, including beetle communities. This study aimed to establish a comprehensive inventory of beetles in the cork oak forest of Larache over two consecutive years (2021 and 2022) by examining beetles associated with different vegetative landscapes using various trapping techniques (pitfall and pan traps) and direct surveys. The results revealed 180 species from 37 families, including 9 endemic species, 33 species newly recorded in Morocco, 13 saproxylic species, and 10 species listed on the Red List (7 saproxylic and 3 coprophagous species). This diversity underscores the ecological importance of the cork oak forest of Larache and highlights the urgent need for conservation measures and the promotion of suitable habitats to protect these insect communities.

Received: 03 December 2024

Accepted: 15 February 2025

Published: 13 April 2025

Subject Editor: Mar Ferrer- Suay

Corresponding author: Amine Samih

E-mail: aminesamih96@gmail.com

DOI: <https://doi.org/10.22034/jesi.45.2.3>**Keywords:** Conservation, Sampling, Diversity, Atlantic coast, Africa

Introduction

The Mediterranean region, encompassing approximately 10% of the world's plant species within just 1.6% of the Earth's land surface (Médail & Quézel, 1997), serves as a vital reservoir of biodiversity. This area supports ecosystems and natural habitats of significant ecological and socio-economic value (Xofis *et al.*, 2024). On its southern shore, forests are indispensable to human societies, providing essential resources such as fuelwood for cooking and heating, fruits, nuts, and medicinal plants that enhance local food security and health (Tsiakiris *et al.*, 2024). Moreover, these forests underpin the livelihoods of many rural communities through activities like resin tapping, cork production, and ecotourism (Fernandes *et al.*, 2024). Beyond their tangible contributions, Mediterranean forests offer critical ecosystem services, including soil erosion control, carbon sequestration, and water regulation functions that are especially crucial in a region highly susceptible to desertification and climatic variability (Fernandes *et al.*, 2024). However, the escalating impacts of global changes, such as climate change, deforestation, land degradation and biodiversity loss, pose severe threats to the sustainability of these ecosystems and their ability to provide essential services to local communities (Bajocco *et al.*, 2012).

In Morocco, the cork oak forests of the Atlantic coastal plain still host various biotopes due to the variability of site conditions and adapted agro-silvo-pastoral systems. Currently, these cork oak woodlands are experiencing severe degradation, which has been exacerbated over the past two decades by the impacts of climate change and significant human pressure. This pressure has intensified due to population growth, the accelerated settlement of communities within forest enclaves and along its edges, and the transformation of traditional land-use practices (Machouri, 2009).

Among the key components of the cork oak forests on the Atlantic coast is the cork oak forest of Larache, which holds significant ecological, economic, and cultural value (Ballouche, 2024). This forest is a crucial habitat

for numerous threatened species (Samih *et al.*, 2024). However, This region is significantly affected by the development of clonal eucalyptus plantations, which deplete local water resources, alter soil composition, and hinder native species regeneration (Peñuelas & Sardans, 2021; Zerga *et al.*, 2021) and modern irrigated agriculture, while boosting crop yields, relies on intensive water use and chemical inputs, contributing to aquifer depletion and soil degradation, further threatening ecosystem sustainability and biodiversity (Benabdelkader *et al.*, 2021). These detrimental practices of deforestation and clearing of cork oak for agriculture and pastoralism disrupt and eliminate habitats for numerous animals, including the insect community. Insects systematically provide essential services such as pollination, organic matter recycling, natural pest regulation, and serving as a food source for many vertebrates (Jactel *et al.*, 2020). In the context of environmental disturbances, the conservation of insects is of paramount importance, as they serve as sensitive bioindicators of the quality of natural ecosystems and the impact of human intervention (Dubucq & Jacquemart, 2020; Sallé *et al.*, 2021).

Due to their bio-ecological characteristics and specialization, beetles have proven to be good indicators for revealing changes in environmental conditions (Davis *et al.*, 2001). Beetles constitute the most diverse and widespread group in the animal kingdom, with approximately 350,000 to 400,000 known species to date, representing about 40% of all insects and 25% of all known animals distribution (Delfosse, 2009; Tronquet, 2021). However, the information on the diversity and distribution of beetles in Morocco is less well known. A recent study in the Maamora forest (Habbaz *et al.*, 2025), which is part of the cork oak forest on the Atlantic coastal plain, provided an inventory of beetles. However, no study has focused on beetle diversity at the scale of the forested areas of the Larache cork oak forest. This study aims to establish a comprehensive inventory of beetles in the Larache cork oak forest. It is therefore necessary to conduct an exhaustive survey of beetles in the cork oak forest of Larache to identify species requiring urgent protection measures.

Materials and methods

Study area

The cork oak forest of Larache, located in the northwest region of Morocco (Fig. 1), forms a geographical unit centered around the Loukkos river plain, bordered to the north and south by the Sahel and R'Mel plateaus (Ballouche, 2013).




Fig. 1. Presentation of the study area and sampled stations.

The climate of the region is Mediterranean, with annual precipitation around 650 mm in Larache, falling mainly between October and May, with 4.5 dry months. Average temperatures are around 18°C, with the coldest month (January) having a minimum average temperature of 5.8°C. Bioclimatically, the region falls within the sub-humid thermomediterranean stage with mild to temperate winters. The prevailing winds are maritime winds from the west to southwest, but in summer, dry and hot continental winds (chergui) are not uncommon, leading to significant evapotranspiration (Ballouche, 2013). The regional vegetation largely consists of cork oak forests, classified as subhumid cork oak forests on sand (Sauvage, 1961). The main forest massif found here is the cork oak forest of Larache, extensively opened up by deforestation and grazing, locally replaced by reforestation with Eucalyptus, Pine and Acacia (Ballouche, 2024).

Description of stations

Three stations were selected within the cork oak forest based on their structure and vegetation (Table 1).

Table 1. Description of the sampled stations.

Station	Locality	Geographic Coordinate	Description
Dense forest (S1)	Larache	35°11'40.9 N 06°07'34.9 W	A dense cork oak forest is distinguished by a closed canopy, varying amounts of deadwood, and an understory primarily composed of Cistaceae and herbaceous plant species (Fig. 2A).
Open forest (S2)	Zouada	35°03'20.2 N 6°01'48.0 W	A degraded cork oak forest bordered by forest tracks and located near cultivated fields. This forest is characterized by a sparse tree cover, often resulting from human activities such as logging and grazing, which reduce tree density and alter habitat structure (Fig. 2B).
Forest edge (S3)	Rissana	35°12'43.1 N 6°01'57.8 W	A flowering edge bordering a dense cork oak forest, characterized by a high diversity of flowering plants. This site benefits from the presence of permanent water resources, including a daya (and aquifer springs (Fig. 2C).



Fig. 2. Sampling stations. A. Dense forest (S1), B. Open forest (S2) and C. Forest edge (S3).

Sampling Methods

Various sampling methods were developed to study the diversity of beetles in the cork oak forest of Larache:

Pan traps

Coloured pan traps (yellow, white, orange, and blue) were installed to capture flower-visiting beetles (Nageleisen *et al.*, 2009). These colored pan traps had a diameter of 15 cm and a height of 13 cm. Each pan trap was filled halfway with water (60 %), detergent (20 %), and salt (20 %) mixture and fixed 1 meter above the ground. These bowls were distributed in four traps per station.

Barber pitfall

Six Barber pitfalls were constructed using plastic cups with a top diameter of 60 mm filled with water (60 %), detergent (20 %), and salt (20 %) and buried flush with the ground to capture species active on the soil surface (Nageleisen & Bouget, 2009).

Malaise trap

A Malaise trap was used to capture specific flying beetles (Ulyshen, 2005; Sheikh *et al.*, 2016), including various families such as Carabidae, Cerambycidae, Chrysomelidae, Curculionidae and Elateridae (Hutcheson, 1990). The Malaise traps were constructed of black and white material and handmade. Each trap measured approximately 1.5 meters in height, 1.2 meters in width, and 1.5 meters in length. The design of the trap is shown in the [fig. 3](#). The traps were installed using stakes driven into the ground and secured with nets. The collection cups contained a solution of 50% ethanol, 5% detergent, and 45% water to preserve the captured specimens. In each station, 11 traps (six Barber pitfall, four Pan traps, and one Malaise trap) were installed and spaced at a distance of 15 to 20 meters apart.

Active trapping

Alongside the passive collection techniques described above, active searching was conducted for the direct capture of individuals, using a sweep net along straight transects (1 m x 100 m) in the sampled sites. We standardized our active searches by making consistent efforts at each site.

Each research session was conducted over a defined period of 8 hours within one day, with a uniform number of 3 researchers. Trap contents were collected every 20 days over 7 months (April-October) over two consecutive years (2021-2022).



Fig. 3. Installation and anchoring of the Malaise trap.

Specimen collection and identification

After each collection session, collected insects were individually placed in plastic vials, labeled, and preserved in 70% ethanol. These specimens were later examined in the laboratory, counted, and identified to species and/or genus level using a binocular microscope and various identification keys (Ferrer, 2009; Alonso, 2007; Baraud, 1992; Ehret, 1990; Delbol & Lempereur, 2014; Jeannel, 1941; Roger *et al.*, 2013; Brendell, 1975; Fadda, 2016; Lillig *et al.*, 2012; Charrier, 2002; Paulian, 1940; Secchi, 2002) and catalogues (Gommy *et al.*, 2011; Trócoli Gracia, 2018; 2019; 2020; 2023 and Carrillo, 2018). For species confirmation, they were compared to the specimens in the museum's collections of the CIRF (Center for Innovation, Research and Training) and the Museum of the Scientific Institute of Rabat (Morocco). The specimens were photographed using an Olympus Stereoscopic Zoom Trinocular Loupe (SZX2-ZB10).

Literature review and species selection

Distribution and endemism

To determine the distribution of captured species at the scale of the Palearctic region and their endemism status in Morocco, we consulted the Catalogues of Palearctic Coleoptera (Löbl & Smetana, 2003; 2004; 2006; 2007; 2008; 2011; 2013).

Rarity and threat status

The rarity and threat status of saproxylic species were determined by consulting the IUCN Red List of Threatened Species, the European Red List of Saproxylic Beetles (Cálix *et al.*, 2018), the Atlas of Threatened Invertebrates of Spain (Critically Endangered Species and Endangered Species) (Verdu & Galante, 2009), the Red List of Invertebrates of Spain (Verdu & Galante, 2006), the Red List of Mediterranean Saproxylic Beetles (García *et al.*, 2018) and the Red List of Coprophagous Beetles (Numa *et al.*, 2020).

Results

To date, this sampling has identified 180 species in this study, belonging to 37 families, among which: 9 are endemic, 33 species are new to Morocco, 13 saproxylic species and 10 are listed in the Red List, including 7 saproxylic species and 3 coprophagous species. The families and species are presented in alphabetical order (Table 2).

[E] endemic species, [N] species new to Morocco, [S] saproxylic species, [R] saproxylic species listed in the Red List of the Mediterranean region, [Rc] coprophagous species listed in the Red List of the Mediterranean region).

Class: Insecta Linnaeus, 1758

Order: Coleoptera Linnaeus, 1758

Family: Anthicidae Latreille, 1819

Hirticollis quadriguttatus (Rossi, 1792)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 28.I.2021, 1 individual, pan traps, 04.VIII.2021, 4 individuals, pitfall trap, 27.IX.2022, 4 individuals, pitfall trap.

Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa, Turkmenistan and Turkey (Chandler *et al.*, 2008).

Omonadus bifasciatus (Rossi, 1792)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 9 individuals, pitfall trap.

Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Algeria, Morocco and Tunisia (Chandler *et al.*, 2008). Ecological insight: The adults are typically found on flowers, where they feed on pollen and nectar. This makes them minor pollinators in the ecosystems they inhabit. The larvae, however, are saprophagous, developing in decaying organic matter, including plant debris or decomposed wood, contributing to nutrient recycling (Telnov, 2010).

Omonadus floralis (Linnaeus, 1758)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps.

Collector name: Samih; Worldwide distribution; Europe, Asia and North Africa (Chandler *et al.*, 2008).

Family: Brentidae Billberg, 1820*Apion frumentarium* (Walton, 1844) [N]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 2 individuals, pan traps.

Collector name; Samih & Maatouf; Worldwide distribution; Europe, Asia and Algeria (Sforzi, 2011).

Hemitrichapion reflexum (Gyllenhal, 1833) [N]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps, 22.VI.2022, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 30.XII.2021, 1 individual, pan traps, 12.V.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and Algeria (Sforzi, 2011).

Family: Bruchidae Latreille, 1802*Bruchidius biguttatus* (Olivier, 1795)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 28.X.2021, 2 individual, pan traps.

Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Anton, 2010; Habbaz *et al.*, 2025).

Bruchidius bimaculatus (Olivier, 1795)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 28.X.2021, 1 individual, pan traps. Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Anton, 2010).

Bruchidius foveolatus (Gyllenhal, 1833)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps, 28.X.2021, 1 individual, pan traps, 12.V.2022, 2 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 1 individual, pan traps, 28.X.2021, 1 individual, pan traps, 30.XII.2021, 1 individual, pan traps. Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Anton, 2010).

Family: Buprestidae Leach, 1815*Acmaeodera adsperula* (Illiger, 1803) [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 2 individuals, pan traps, 15.IX.2021, 1 individual, pan traps, 22.VI.2022, 1 individual, Pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 3 individuals, pitfall trap, 22.VI.2022, 1 individual, pitfall trap.

Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia, Morocco, Algeria and Tunisia (Villemant & Fraval, 1993; Volkovitsh, 2006; Habbaz *et al.*, 2025).

Ecological insight

This species is commonly associated with woody plants, especially oak trees, where its larvae develop within the decaying wood, contributing to nutrient cycling within forest ecosystems. This behavior is characteristic of many wood-boring beetles, including those in the genus *Acmaeodera* (Alizadeh *et al.*, 2024).

Acmaeodera discoida (Fabricius, 1787)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 1 individual, pitfall trap; Collector name: Samih; Worldwide distribution: Europe, North Africa, Syria and Israël (Volkovitsh, 2006; Habbaz *et al.*, 2025).

Acmaeodera lanuginosa (Gyllenhal, 1817)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 3 individuals, pan traps, 01.VII.2021, 1 individual, pitfall trap, 27.V.2021, 5 individuals, pan traps, 04.VIII.2021, 1 individual, pan traps, 29.IV.2022, 3 individuals, pan traps, 12.V.2022, 1 individual, pitfall trap, 01.VI.2022, 6 individuals, pitfall trap, 22.VI.2022, 1 individual, pan traps. Collector name Samih: Maatouf & Habbaz; Worldwide distribution: Europe, Morocco, Algeria, Tunisia, Sinai and Israel (Volkovitsh, 2006; Habbaz *et al.*, 2025).

Agrilus biguttatus (Fabricius, 1777) [S]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, malaise trap. Collector name: Samih; Worldwide distribution Europe, Iran, Syria, Algeria and Morocco (Jendek, 2006; Habbaz *et al.*, 2025).

Agrilus graminis mamorensis (Théry, 1930) [E] [S]

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 3 individuals, pan traps; Collector name; Samih; Worldwide distribution: Morocco (Jendek, 2006; Habbaz *et al.*, 2025).

Anthaxia millefolii polychloros (Abeille de Perrin, 1894)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 2 individuals, pan traps, 01.VI.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps, 27.V.2021, 3 individuals, pan traps, 04.VIII.2021, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe and North Africa (Bily, 2006; Habbaz *et al.*, 2025).

Anthaxia scutellaris atlasica (Théry, 1930) (Fig. 4B)

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 7 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Algeria, Tunisia and Morocco (Bily, 2006).

Anthaxia umbellatarum umbellatarum (Fabricius, 1787) (Fig. 4A)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 22.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps, 27.V.2021, 1 individual, pan traps, 01.VII.2021, 2 individuals, pan traps, 15.IX.2021, 3 individuals, pitfall trap, 01.VI.2022, 1 individual,

pan traps, 21.VII.2022, 2 individuals, pan traps, 27.IX.2022, 3 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia and North Africa (Bily, 2006; Habbaz *et al.*, 2025).

Habroloma triangulare (Lacordaire, 1835)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 2 individuals, pan traps, 27.IX.2022, 2 individuals, pan traps; Collector name Samih; Worldwide distribution: Spain, France, Italy, Portugal, Algeria, Tunisia and Morocco (Kuban, 2006; Habbaz *et al.*, 2025).

Family: Carabidae Latreille, 1802

Acupalpus brunnipes (Sturm, 1825)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 1 individual, pan traps, 22.VI.2022, 1 individual pan traps; Collector name: Samih; Worldwide distribution: Europe, Algeria, Tunisia, Morocco and Turkey (Jaeger & Kataev, 2003).

Agonum emarginatum (Gyllenhal, 1827) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 4 individuals, pan traps, 22.VI.2022, 4 individuals, pan traps; Collector name Samih & Maatouf; Worldwide distribution Europe (Bousquet, 2003a).

Bembidion biguttatum (Fabricius, 1779) (Fig. 4F) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and Canary Islands (Marggi *et al.*, 2003). Worldwide distribution: The adults of *Bembidion* genus, as predators of small arthropods, contribute to maintaining the balance of invertebrate communities by regulating the populations of certain species (Ekschmitt *et al.*, 1997; Staudacher *et al.*, 2024).

Bembidion bipunctatum laevifrons (Schaufuss, 1882)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 1 individual, pan traps; Collector name Maatouf; Worldwide distribution Spain, Portugal and Morocco (Marggi *et al.*, 2003).

Bradycellus verbasci (Duftschmid, 1812)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.IX.2022, 2 individuals, pan traps; Collector name Maatouf; Worldwide distribution Europe, Asia and North Africa (Bousquet, 2003a).

Carterus rotundicollis (Rambur, 1837)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Algeria, Tunisia and Morocco (İto, 2003).

Cincidela Moroccana (Fabricius, 1801) (Fig. 4E)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 28.10.2021, 1 individual, pitfall trap; Collector name Samih; Worldwide distribution: Portugal, Spain, Tunisia and Morocco (Puchkov & Matalin, 2003; Habbaz *et al.*, 2025).

Cymindis lineola (L. Dufour, 1820) (Fig. 4I) [N]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pitfall trap.

Rissana, Open forest, 35°10'48.0 N 5°58'40.1 W, 28.X.2021, 1 individual, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe and Tunisia (Kabak, 2003).

Cymindis platicollis (Say, 1823) [N]

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 07.VI.2021, 2 individuals, pan traps; Collector name Samih; Worldwide distribution: Canada and Mexico (Arce-Pérez *et al.*, 2018).

Harpalus attenuatus (Stephens, 1828)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 1 individual, pan traps, 15.IX.2021, 3 individuals, pitfall trap, 28.X.2021, 5 individuals, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Kataev *et al.*, 2003; Habbaz *et al.*, 2025).

Microlestes abeillei brisouti (Holdhaus, 1912)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 3 individuals, pan traps, 01.VI.2022, 1 individual, pan traps; Collector name Samih & Maatouf; Worldwide distribution North Africa (Kabak, 2003; Habbaz *et al.*, 2025).

Ophonus ardosiacus (Lutshnik, 1922)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps, 12.V.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa and Turkey (Kataev *et al.*, 2003).

Ophonus subquadratus (Dejean, 1829) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 8 individuals, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia, Algeria and Libya (Kataev *et al.*, 2003).

Paradromius linearis (Olivier, 1795) (Fig. 4D)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 28.10.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Kabak, 2003; Habbaz *et al.*, 2025).

Philorhizus notatus (Stephens, 1827) [N]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps; Collector name Samih; Worldwide distribution: Europe and Asia (Kabak, 2003).

Pterostichus elongatus (Duftschmid, 1812)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 28.X.2021, 3 individuals, pitfall trap, 01.VI.2022, 5 individuals, pitfall trap, 22.VI.2022, 12 individuals, pitfall trap, 27.IX.2022, 2 individuals, pitfall trap, 30.12.2022, 5 individuals, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 07.VII.2021, 5 individuals, pitfall trap, 28.X.2021, 20 individuals, pitfall trap, 30.XII.2021, 6 individuals, pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 30.XII.2021, 3 individuals, pitfall trap, 12.V.2022, 5 individuals, pitfall trap, 22.VI.2022, 1 individuals, pitfall trap; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Morocco, Afghanistan, Kazakhstan and Turkey (Bousquet, 2003b; Habbaz *et al.*, 2025).

Scarites terricola terricola (Bonelli, 1813) (Fig. 4C)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pitfall trap, 01.VI.2022, 4 individuals, pitfall trap, 22.VI.2022, 2 individuals, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Balkenohl, 2003; Habbaz *et al.*, 2025).

Singilis soror soror (Rambur, 1837)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 15.IX.2021, individual, pan traps, 21.VII.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Spain, Algeria and Morocco (Kabak, 2003).

Stenolophus abdominalis abdominalis (Gene, 1836)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 2 individuals, pan traps, 12.V.2022, 5 individuals, pan traps, 21.VII.2022, 7 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: France, Italy, Portugal, Spain, North Africa and Saudi Arabia (Jaeger & Kataev, 2003).

Stenolophus teutonius (Schränk, 1781)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.IX.2022, 6 individuals, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Jaeger & Kataev, 2003).

Syntomus foveatus (Geoffroy, 1785)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 3 individuals, pan traps; Collector name Samih; Worldwide distribution: Europe, Asia and North Africa (Kabak, 2003; Habbaz *et al.*, 2025).

Family: Cerambycidae Latreille, 1802

Agapanthia irrorata (Fabricius, 1787)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, malaise trap; Collector name: Maatouf; Worldwide distribution: France, Italy, Spain, Portugal, Algeria, Morocco and Tunisia (Sama, 2010).

Alocerus moesiacus (Frivaldszky, 1837) (Fig. 4J)

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 1 individual, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Algeria, Morocco and Tunisia (Sama, 2010).

Certallum ebulinum (Linné, 1767)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 2 individuals, malaise trap; Collector name: Maatouf; Worldwide distribution; Europe, Asia and North Africa (Sama, 2010).

Chlorophorus favieri (Farimaire, 1873) (Fig. 4G) [E] [R]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Morocco (Sama, 2010); Ecological insight: The larvae of *C. favieri* are saproxylic, develop in the dying branches of oaks (Sama, 1987).

Phytoecia coerulea (Scopoli, 1763)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, malaise trap; Collector name: Samih; Worldwide distribution: Europe, Asia, Algeria, Morocco and Tunisia (Sama, 2010).

Oxypleurus nodieri (Mulsant, 1839)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, North Africa and Turkey (Sama, 2010).

Stenurella approximans (Rosenhauer, 1856)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps, 01.VI.2022, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.V.2022, 2 individuals, malaise trap, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Portugal, Spain and Morocco (Sama, 2010).

Stictoleptura fontenayi (Mulsant & Rey, 1839)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Algeria, Morocco and Tunisia (Sama, 2010).

Trichoferus ilicis (Sama, 1987) [E] [R]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 28.X.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 5 individuals, pan traps, 28.X.2021, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Morocco (Sama, 2010; Habbaz et al., 2025).

Family: Chrysomelidae Latreille, 1802*Aphthona euphorbiae* (Schrank, 1781)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Kippenberg, 2010).

Cassida vittata (Villers, 1789)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 2 individuals, pan traps, 01.VII.2021, 11 individuals, pan traps, 30.XII.2021, 4 individuals, pan traps, 01.VI.2022, 2 individuals, pan traps, 22.VI.2022, 4 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Borowiec & Sekerka, 2010).

Chrysolina bankii (Fabricius, 1775)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe and North Africa (Kippenberg, 2010).

Chrysolina diluta (Germar, 1823) [N]**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 30.XII.2021, 1 individual, pitfall trap; Collector name: Maatouf; Worldwide distribution: France, Spain and Portugal (Kippenberg, 2010).

Cryptocephalus fulvus (Goeze, 1777) [N]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 1 individual, pan traps; Collector name: Samih & Habbaz; Worldwide distribution: Europe, Kazakhstan, Tajikistan and Russia (Lopatin *et al.*, 2010).

Cryptocephalus numidicus (Bourdonné, 1876)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, direct surveys; Collector name: Habbaz; Worldwide distribution: Algeria, Morocco and Tunisia (Lopatin *et al.*, 2010). Ecological insight: Genus *Cryptocephalus* have unique ecological habits: larvae feed on dead vegetable material and case-bearer females envelop surface of eggs with their excrement (Kang & Lee, 2017).

Longitarsus aeneus (Kutschera, 1862)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 30.XII.2021, 2 individuals, pan traps, 01.VI.2022, 2 individuals, malaise trap, 22.VI.2022, 2 individuals, malaise trap, 27.IX.2022, 1 individual, pan traps. Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, North Africa, Syria, Yemen and Jordan (Doberl, 2010).

Longitarsus ochroleucus (Marshall, 1802)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 04.VIII.2021, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 13 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa, China and Turkey (Doberl, 2010).

Oulema melanopus (Linnaeus, 1758)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Schmitt, 2010).

Psylliodes cuprea (Koch, 1803)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps, 27.IX.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Algeria, Morocco and Tunisia (Doberl, 2010).

Family: Coccinellidae Latreille, 1807*Adalia decempunctata* (Linnaeus, 1758)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Kovar, 2007; Habbaz *et al.*, 2025).

Chilocorus bipustulatus (Linnaeus, 1758)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Kovar, 2007).

Coccinella septempunctata (Linnaeus, 1758)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Italy and North Africa (Kovar, 2007).

Oenopia conglobata (Linnaeus, 1758) (Fig. 4L)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 28.X.2021, 3 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia (Kovar, 2007) and Morocco (Benyahia *et al.*, 2016; Habbaz *et al.*, 2025); Ecological insight: Both the adult and larval stages of *O. conglobata* are predatory, primarily feeding on aphids, which are their main food source. They are effective natural pest controllers in ecosystems where aphids are prevalent (Lumbierres *et al.*, 2018).

Oenopia lyncea lyncea (Olivier, 1808)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: France, Italy, Portugal, Spain, Algeria, Morocco and Tunisia (Kovar, 2007; Habbaz *et al.*, 2025).

Platynaspis luteorubra (Goeze, 1777)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Kovar, 2007; Habbaz *et al.*, 2025).

Rhyzobius litura (Fabricius, 1787)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pan traps, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa and Turkey (Kovar, 2007; Habbaz *et al.*, 2025).

Rhyzobius lophantae (Blaisdell, 1892)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa, Iran, Inde and Turkey (Kovar, 2007).

Rodolia cardinalis (Mulsant, 1850) (Fig. 4K)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Kovar, 2007; Habbaz *et al.*, 2025).

Scymnus abiandis (Paykull, 1798) [N]**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe and Asia (Kovar & Smetana, 2007).

Scymnus apandzi (Mulsant, 1846) [N]**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 3 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe and Asia (Kovar, 2007).

Scymnus impexus (Mulsant, 1850) [N]**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 4 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe and China (Kovar, 2007).

Scymnus interruptus (Goeze, 1777)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 28.X.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Kovar, 2007).

Scymnus subvillosus (Goeze, 1777)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Kovar, 2007; Habbaz *et al.*, 2025).

Scymnus suturalis (Westman, 1795) [N]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 21.VII.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia, Algeria, Madeira Islands and Tunisia (Kovar, 2007).

Standhorus punctillum (Weise, 1891)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 04.VIII.2021, 1 individual, pan traps, 12.V.2022, 2 individuals, pan traps, 01.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Egypt and Morocco (Kovar, 2007).

Family: Corylophidae LeConte, 1852*Arthrolips convexiuscula* (Motschulsky, 1849)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 3 individuals, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Bowstead, 2007).

Family: Curculionidae Latreille, 1802*Brachyderes incanus* (Linné, 1758)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 2 individuals, pan traps, 12.V.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 28.X.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Japan (Pellandier, 2013) and Morocco (Arahou, 2008).

Brachyderes pubescens (Boheman, 1833)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.IX.2022, 3 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Spain, Portugal, France, Italy, Algeria and Morocco (Pellandier, 2013; Habbaz et al., 2025).

***Brachytemnus porcatus* (Germar, 1823)**

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps, 01.VI.2022, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Algeria, Morocco, Lebanon, Syria, Turkey and Cyprus (Hlavac & Maughan, 2013).




Fig. 4. The beetles associated with the Larache cork oak forest. A. *Anthaxia umbellatarum umbellatarum* (Fabricius, 1787), B. *Anthaxia scutellaris atlasica* (Théry, 1930), C. *Scarites terricola terricola* (Bonelli, 1813), D. *Paradromius linearis* (Olivier, 1795), E. *Cicindela maroccana* Fabricius, 1801, F. *Bembidion biguttatum* (Fabricius, 1779), G. *Chlorophorus favieri* (Farimaire, 1873), H. *Oedemera barbara* (Fabricius, 1792), I. *Cymindis lineola* (L. Dufour, 1820), J. *Alocerus moesiacus* (Frivaldszky, 1837), K. *Rodolia cardinalis* (Mulsant, 1850), L. *Oenopia conglobata* (Linnaeus, 1758) (photos by A. Samih and N. Maatouf).

Ceutorhynchus pallidactylus (Marsham, 1802)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.IX.2022, 1 individual, pan traps; Collector name Maatouf; Worldwide distribution Europe, Asia, and North Africa (Colonnelli, 2013).

Coeliodes ruber (T.Marsham, 1802)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 2 individuals, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Algeria, Morocco, Syria and Israel (Colonnelli, 2013; Habbaz *et al.*, 2025).

Lixus juncii (Boheman, 1835) [N]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 2 individuals, pan traps, 27.IX.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Algeria, Madeira Islands, Egypt, Syria, Iraq, Jordan, Cyprus and Turkey (Gultekin & Fremuth, 2013).

Microplontus rugulosus (J.F.W.Herbst, 1795)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.IX.2022, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Caldara, 2013).

Orchestes irroratus maroccanus (Roudier, 1954)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps, 01.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Algeria, Morocco and Tunisia (Villemant & Fraval, 1993; Caldara, 2013).

Sitona callosus (Gyllenhal, 1834) [N]**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 30.XII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe and Asia (Velazquez de castro, 2013).

Sitona lineatus (Linnaeus, 1758) [N]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 2 individuals, pan traps, 01.VI.2022, 2 individuals, pan traps, 21.VII.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe North Africa and Asia (Velazquez de castro, 2013).

Sitona lineellus (Bonsdorff, 1785) [N]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 2 individuals, pan traps; Collector name: Samih; Worldwide distribution: Europe and Asia (Velazquez de castro, 2013).

Sitona longulus (Gyllenhal, 1834) [N]**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 15.IX.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps; Collector name Samih & Maatouf; Worldwide distribution Europe and Asia (Velazquez de castro, 2013).

Tychius cuprifer (Panzer, 1799)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Algeria, Morocco and Asia (Caldara, 2013).

Tychius pusillus (Germar, 1842)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 2 individuals, pan traps, 01.VI.2022, 1 individual, pan traps, 22.VI.2022, 1 individual, pan traps, 22.VI.2022, 2 individuals, malaise trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Algeria and Morocco (Caldara, 2013).

Family: Dasytidae Laporte, 1840*Dasytes nigroaeneus* (Küster, 1850)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 38 individuals, pan traps, 27.V.2021, 8 individuals, pan traps, 01.VII.2021, 5 individuals, pan traps, 12.V.2022, 24 individuals, pan traps, 12.V.2022, 2 individuals, malaise trap, 01.VI.2022, 12 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 12 individuals, pan traps, 27.V.2021, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 20.IV.2021, 31 individuals, pan traps, 27.V.2021, 46 individuals, pan traps, 01.VII.2021, 12 individuals, pan traps, 12.V.2022, 95 individuals, pan traps, 01.VI.2022, 41 individuals, pan traps, 22.VI.2022, 12 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Syria, Algeria and Morocco (Mayor, 2007a; Habbaz et al., 2025).

Dasytes terminalis (Jacquelin du Val, 1863)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 8 individuals, pan traps, 27.V.2021, 8 individuals, pan traps, 01.VII.2021, 2 individuals, pan traps, 12.V.2022, 10 individuals, pan traps, 12.V.2022, 4 individuals, malaise trap, 01.VI.2022, 50 individuals, pan traps, 22.VI.2022, 8 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 2 individuals, pan traps, 01.VII.2021, 1 individual, pan traps, 04.VIII.2021, 1 individual, pitfall trap, 30.XII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 12 individuals, pan traps, 27.V.2021, 9 individuals, pan traps, 01.VII.2021, 7 individuals, pan traps, 12.V.2022, 48 individuals, pan traps, 01.VI.2022, 18 individuals, pan traps, 01.VI.2022, 4 individuals, malaise trap, 22.VI.2022, 3 individuals, pan traps, 21.VII.2022, 1 individual, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: France, Portugal, Spain, Algeria and Morocco (Mayor, 2007a; Bellifa & Roussand, 2021; Habbaz et al., 2025).

Family: Dermestidae Latreille, 1804*Anthrenus flavipes* (LeConte, 1854)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 3 individuals, pan traps, 21.VII.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Egypt, Tunisia and Morocco (Hava, 2007b).

Anthrenus fuscus (Olivier, 1790)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 1 indivdu, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 2 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps, 21.VII.2022, 2 individuals, pan traps; Collector name Samih & Maatouf; Worldwide distribution Europe and Morocco (Hava, 2007b).

Anthrenus museorum (Linnaeus, 1761)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Hava, 2007b; Habbaz *et al.*, 2025).

Anthrenus pimpinellae (Fabricius, 1775)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia, Egypt and Morocco (Hava, 2007b; Habbaz *et al.*, 2025).

Attagenus bifasciatus (Fabricius, 1787)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 7 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 6 individuals, pan traps, 21.VII.2021, 6 individuals, pan traps, 12.V.2022, 6 individuals, pan traps, 01.VI.2022, 19 individuals, pan traps, 22.VI.2022, 6 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia and North Africa (Hava, 2007b).

Dermestes frischii (Kugelann, 1792)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 2 individuals, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 2 individuals, pitfall trap, 04.VIII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 4 individuals, pitfall trap, 21.VII.2022, 2 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia and North Africa (Hava, 2007a; Habbaz *et al.*, 2025); Ecological insight Is a necrophagous species, helps accelerate the breakdown of dead organisms, returning valuable nutrients to the soil (Farag *et al.*, 2021).

Orphilus niger (Rossi, 1790)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Uzbekistan, Tajikistan and North Africa (Hava, 2007a).

Trogoderma granarium (Everts, 1898)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 2 individuals, pan traps, 12.V.2022, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Egypt, Tunisia and Morocco (Hava, 2007a).

Family: Dryopidae Billberg, 1820

Dryops luridus (Erichson, 1847) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 30.XII.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Turkey and North Africa except Morocco (Kodada & Jäch, 2006).

Family: Elateridae Leach, 1815*Cardiophorus rufipes* (Buysson, 1902) [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 2 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 2 individuals, pan traps, 27.V.2021, 3 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 3 individuals, pan traps, 01.VI.2022, 3 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia and North Africa (Cate, 2007) and Morocco (Habbaz et al., 2025).

Drasterius bimaculatus (Rossi, 1790) [S]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps, 30.XII.2021, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa, Asia and Morocco (Cate, 2007).

Family: Erotylidae Latreille, 1802*Triplax lacordairei* (Crotch, 1870) (Fig. 5I) [R]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Algeria and Morocco (Wegrzynowicz, 2007a).

Family: Geotrupidae Latreille, 1802*Thorectes distinctus* (Marseul, 1878) (Fig. 5E)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 11 individuals, pitfall trap, 12.V.2022, 20 individuals, pitfall trap, 01.VI.2022, 24 individuals, pitfall trap, 22.VI.2022, 13 individuals, pitfall trap, 21.VII.2022, 5 individuals, pitfall trap, 27.IX.2022, 1 individual, pitfall trap, 30.XII.2022, 57 individuals, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 15.IX.2021, 2 individuals, pitfall trap, 28.X.2021, 5 individuals, pitfall trap, 30.XII.2021, 58 individuals, pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 11 individuals, pitfall trap, 01.VI.2022, 8 individuals, pitfall trap; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Algeria and Morocco (Löbl et al., 2006; Habbaz et al., 2025).

Typhaeus typhoeus (Linnaeus, 1758)**Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 30.XII.2021, 1 individual, pitfall trap; Collector name: Maatouf; Worldwide distribution: Europe and Morocco (Löbl et al., 2006; Habbaz et al., 2025).

Family: Glaphyridae Macleay, 1819*Anthypna meles* (Fabricius, 1792)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 22.VI.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Algeria, Tunisia and Morocco (Nikodym & Bezdek, 2006; Habbaz et al., 2025).

Eulasia goudoti (Laporte, 1840) [E]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 4 individuals, pan traps, 01.VI.2022, 9 individuals, pan traps, 01.VI.2022, 3 individuals, pitfall trap, 01.VII.2022, 3 individuals, pan traps. Collector name: Samih

& Maatouf; Worldwide distribution: Morocco (Nikodym & Bezdek, 2006; Habbaz *et al.*, 2025); Ecological insight: Characterized by its close interactions with flowering plants, it feeds on pollen and nectar, thus playing a role in pollination (Samih *et al.*, 2024).

Family: Histeridae Gyllenhal, 1808

Hypocaccus rugiceps (Duftschmid, 1805)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 2 individuals, pan traps, 21.VII.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe and China (Mazur, 2004; Habbaz *et al.*, 2025).

Saprinus proximus simillimus (Wollaston, 1865)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 8 individuals, pitfall trap, 01.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: North Africa (Mazur, 2004) and Morocco (Habbaz *et al.*, 2025).

Family: Hydraenidae Mulsant, 1844

Ochthebius bicolon (Germar, 1823) [N]

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 28.X.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe (Jäch, 2004).

Family: Hydrophilidae Latreille, 1802

Cercyon obsolandus (Gyllenhal, 1808) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Algeria and Canary Islands (Hansen, 2004).

Sphaeridium scarabaeoides (Linnaeus, 1758) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2021, 1 individual, pan traps. 21.VII.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and Tunisia (Hansen, 2004).

Family: Laemophloeidae Ganglbauer, 1899

Laemophloeus monilis (Fabricius, 1787) [S]

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Uzbekistan, Tunisia, Algeria (Wegrzynowicz, 2007b) and Morocco (Benyahia, 2016; Habbaz *et al.*, 2025).

Laemophloeus muticus (Fabricius, 1781) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe and Russie (Wegrzynowicz, 2007b).

Family: Latridiidae Erichson, 1842

Corticarina curta (Wollaston, 1854)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia and North Africa (Johnson, 2007; Habbaz *et al.*, 2025).

Enicmus transversus* (Olivier, 1790)*Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Afghanistan, China, Israel, Jordan, Syria and North Africa (Johnson, 2007).

Stephostethus productus* (Rosenhauer, 1856)*Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pitfall trap, 15.IX.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: France, Italy, Portugal, Spain and North Africa (Johnson, 2007).

Family: Meloidae Gyllenhaal, 1810***Berberomeloe majalis* (Linnaeus, 1758) (Fig. 5A)****Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 04.VIII.2021, 1 individual, direct surveys; Collector name: Samih; Worldwide distribution: France, Portugal, Spain, Algeria, Tunisia and Morocco (Bologna, 2008).

Croscherichia paykulli* (Billberg, 1813)*Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 15.IX.2021, 1 individual, pitfall trap; Collector name: Maatouf; Worldwide distribution: Algeria, Tunisia, Libya and Morocco (Bologna, 2008).

Hycleus rufipalpis* (Escalera, 1909)*Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 3 individuals, pitfall trap, 04.VIII.2021, 1 individual, pitfall trap, 15.IX.2021, 1 individual, pitfall trap, 28.X.2021, 2 individuals, pan traps, 22.VI.2022, 1 individual, pan traps, 21.VII.2022, 1 individual, pitfall trap, 21.VII.2022, 2 individuals, pan traps, 27.IX.2022, 9 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Canary Islands and Morocco (Bologna, 2008; Habbaz et al., 2025).

Mylabris variabilis* (Pallas, 1781) (Fig. 5D)*Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia (Bologna, 2008) and Morocco (Pan et al., 2014).

Family: Melolonthidae Leach, 1819***Euserica mamorensis* (Baraud, 1965) [E]****Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 20.IV.2021, 1 individual, pan traps, 01.VII.2021, 1 individual, pan traps, 01.VI.2022, 3 individuals, pan traps, 22.VI.2022, 3 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 1 individual, pan traps, 01.VII.2021, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 1 individual, pan traps, 01.VII.2021, 2 individuals, pan traps, 04.VIII.2021, 1 individual, pan traps, 04.VIII.2021, 1 individual, pitfall trap, 12.V.2022, 1 individual, malaise trap, 22.VI.2022, 1 individual, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Morocco (Bezdek, 2006; Habbaz et al., 2025).

Hoplia africana* (Escalera, 1914)*Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 7 individuals, pan traps, 27.V.2021, 12 individuals, pan traps, 12.V.2022, 4 individuals, pan traps, 22.VI.2022, 7 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Algeria, Morocco and Tunisia (Smetana, 2006).

Hoplia bilineata (Fabricius, 1801)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 5 individuals, pan traps; Collector name: Maatouf; Worldwide distribution: Spain, Algeria, Morocco and Tunisia (Smetana, 2006).

Hoplia philanthus (Fuessly, 1775)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps.

Forest edge, 27.V.2021, 3 individuals, pan traps, 01.VII.2021, 5 individuals, pan traps, 12.V.2022, 3 individuals, pan traps, 21.VII.2022, 3 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Spain and Morocco (Smetana, 2006).

Sphodroxia Moroccana (Ley, 1923) [E]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 15.IX.2021, 1 individual, pitfall trap, 27.IX.2022, 3 individuals, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Morocco (Bezdek, 2006; Habbaz *et al.*, 2025).

Family: Melyridae Leach, 1815

Aplocnemus virens (Suffrian, 1843) [N]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 2 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps, 21.VII.2022, 2 individuals, pan traps. Collector name: Samih & Maatouf; Worldwide distribution: Europe (Mayor, 2007b).

Colotes javeti (Jacquelin du Val, 1852)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Spain, France, Italy, Portugal, Egypt, Tunisia, Algeria and Morocco (Mayor, 2007; Habbaz *et al.*, 2025).

Psilothrix viridicoerulea (Geoffroy, 1758)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, North Africa and Turkey (Mayor, 2007; Amokrane *et al.*, 2020).

Troglops furcatus (Perrin, 1885)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps, 01.VI.2022, 5 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 4 individuals pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Portugal, Spain, Algeria and Morocco (Mayor, 2007b)

Family: Mordellidae Latreille, 1802***Mediimorda bipunctata* (Germar, 1827)****Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 5 individuals, pan traps, 04.VIII.2021, 19 individuals, pan traps, 15.IX.2021, 17 individuals, pitfall trap, 21.VII.2022, 1 individual, pan traps, 27.IX.2022, 2 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution Europe, Asia, North Africa (Horak, 2008) and Morocco (Habbaz et al., 2025).

Mordella aculeata* (Linnaeus, 1758)*Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps, 01.VII.2021, 1 individual, pan traps, 01.VI.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 3 individuals, pan traps, 21.VII.2022, 2 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia (Horak, 2008), Austria, Belgium, Bosnia, Herzegovina, Bulgaria, Croatia, Czech Republic, Denmark, France, Germany, Greece, Hungary, Italy, Albania, Macedonia, Poland, Portugal, Romania, Serbia, Montenegro, Slovakia, Slovenia, Spain, Sweden, Switzerland, Syria, Turkey, Georgia, Ukraine, Belarus, Russia, Kazakhstan, China, Taiwan, Japan, South Korea (Ruzzier et al., 2017) and Morocco (Habbaz et al., 2025).

Variimorda villosa* (Schrank von Paula, 1781) [N]*Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 21.VII.2022, 1 individual, pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 6 individuals, pan traps, 04.VIII.2021, 40 individuals, pan traps, 19.IX.2021, 9 individuals, pan traps, 28.X.2021, 6 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps, 21.VII.2022, 60 individuals, pan traps, 27.IX.2022, 9 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe and Asia (Horak, 2008).

Family: Nitidulidae Latreille, 1802***Carpophilus hemipterus* (Linnaeus, 1758)****Material examined**

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 28.X.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Algeria, Canary Islands, Egypt, Libya, Madeira Island, Tunisia (Jelinek & Audisio, 2007) and Morocco (Chavanon, 2018; Habbaz et al., 2025).

Epuraea latipes* (Grouvelle, 1896)*Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 2 individuals, pan traps; Collector name: Maatouf; Worldwide distribution: Algeria, Morocco and Turkey (Jelinek & Audisio, 2007).

Meligethes aeneus* (Fabricius, 1775)*Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 5 individuals, pan traps, 27.V.2021, 1 individual, pan traps, 12.V.2022, 1 individual, pan traps, 21.VII.2022, 2 individuals, pan traps. Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 3 individuals, pan traps, 01.VII.2021, 2 individuals, pan traps, 04.VIII.2021, 3 individuals, pitfall trap, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia and North Africa (Jelinek & Audisio, 2007).

Meligethes viridescens (Fabricius, 1787)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 28.X.2021, 1 individual, pan traps, 12.V.2022, 2 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 29.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 2 individuals, pan traps, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia, Algeria, Morocco and Tunisia (Jelinek & Audisio, 2007).

Family: Oedemeridae Latreille, 1810*Chrysanthia viridissima* (Linnaeus, 1758) [N]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 22 individuals, pan traps, 27.V.2021, 98 individuals, pan traps, 01.VII.2021, 2 individuals, pan traps, 12.V.2022, 2 individuals, pan traps, 01.VI.2022, 7 individuals, pan traps, 22.VI.2022, 1 individual, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe and Russia (Švihla, 2008).

Oedemera barbara (Fabricius, 1792) (Fig. 4H) [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 8 individuals, pan traps, 01.VII.2021, 3 individuals, pan traps, 29.IV.2022, 2 individuals, pan traps, 01.VI.2022, 10 individuals, pan traps, 22.VI.2022, 3 individuals, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 19 individuals, pan traps, 27.V.2022, 4 individuals, pan traps, 01.VI.2022, 2 individuals, pitfall trap, 22.VI.2022, 5 individuals, pan traps, 01.VII.2022, 22 individuals, pan traps, 21.VII.2022, 1 individual, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, North Africa, Turkey (Švihla, 2008) and Morocco (Habbaz *et al.*, 2025); Ecological insight Is considered a pollinator, with adult beetles visiting flowers to feed on pollen and nectar (Samih *et al.*, 2024).

Oedemera marmorata (Erichson, 1841)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2022, 5 individuals, pan traps, 01.VI.2022, 29 individuals, pan traps, 22.VI.2022, 4 individuals, pan traps, 01.VII.2022, 5 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Portugal, Spain, Algeria, Morocco and Tunisia (Švihla, 2008).

Family: Phalacridae Leach, 1815*Olibrus pygmaeus* (Sturm, 1807)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 04.VIII.2021, 1 individual, pan traps, 01.VI.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IV.2021, 3 individuals, pan traps, 15.IX.2021, 2 individuals, pan traps, 22.VI.2022, 1 individuals, pan traps, 21.VII.2022, 4 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, North Africa, Syria and Turkey (Švec, 2008).

Phalacrus coruscus (Panzer, 1797)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 29.IX.2022, 2 individuals, pan traps; Collector name: Samih; Worldwide distribution: Europe, North Africa and Asia (Švec, 2008).

Family: Ptinidae Latreille, 1802*Dignomus irroratus* (Kiesenwandter, 1851) [N]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 29.IV.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 1 individual, pan traps, 01.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe (Borowski, 2007).

Stegobium paniceum (Linnaeus, 1758) [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 04.VIII.2021, 1 individual, pan traps, 28.X.2021, 1 individual, pan traps, 01.VI.2022, 1 individual, pan traps, 22.VI.2022, 1 individual, pan traps.

Rissana, Forest edge 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 4 individuals, pan traps, 15.IX.2021, 4 individuals, pan traps, 22.VI.2022, 4 individuals, pan traps, 01.VII.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, North Africa (Borowski, 2007; Habbaz *et al.*, 2025).

Family: Scarabaeidae Latreille, 1802*Anisoplia baetica* (Erichson, 1847) [N]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 2 individuals, pan traps, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Portugal and Spain (Smetana, 2006).

Anthoplia floricola (Fabricius 1787)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Portugal, Spain, Algeria, Morocco, Tunisia and Libya (Smetana, 2006).

Aphodius diecki (Harold, 1870)**Material examined**

Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 28.X.2021, 1 individual, pitfall trap; Collector name: Samih; Worldwide distribution: Europe, Asia, Algeria, Tunisia and Morocco (Dellacasa & Dellacasa, 2006).

Blitopertha lineata (Fabricius, 1798)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 22.VI.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Spain, Algeria, Morocco and Tunisia (Smetana, 2006).

Gymnopleurus flagellatus (Fabricius, 1787)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 1 individual, pitfall trap; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa (Löbl *et al.*, 2006).

Gymnopleurus sturmii (MacLeay, 1821) [Rc]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 1 individual, pitfall trap; Collector name: Samih; Worldwide distribution: Europe, Jordan, Israel, Syria, Turkey and North Africa (Löbl *et al.*, 2006).

Onthophagus maki (Illiger, 1803) (Fig. 5F) [Rc]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 4 individuals, pitfall trap, 01.VI.2022, 4 individuals, pitfall trap, 22.VI.2022, 2 individuals, pitfall trap; Collector name: Samih & Maatouf; Worldwide

distribution Spain, France, Portugal, Italy, North Africa (Löbl *et al.*, 2006a) and Morocco (Habbaz *et al.*, 2025); Ecological insight: as a dung beetle, *Onthophagus* contributes to nutrient cycling by breaking down dung into smaller pieces, enhancing nutrient availability for plants. The process of dung burial helps incorporate organic matter into the soil (Ramli *et al.*, 2019).




Fig. 5. The beetles associated with the Larache cork oak forest. A. *Berberomeloe majalis* (Linnaeus, 1758), B. *Scarabaeus sacer* (Linnaeus, 1758), C. *Scarabaeus cicatricosus* (P.H.Lucas, 1846), D. *Mylabris variabilis* (Pallas, 1781), E. *Thorectes distinctus* (Marseul, 1878), F. *Onthophagus maki* (Illiger, 1803), G. *Trichius zonatus* (Germar, 1831), H. *Endophloeus markovichianus* (Piller & Mitterpacher, 1783), I. *Triplax lacordairei* (Crotch, 1870), J. *Cossyphus hoffmanseggii* (Herbst, 1797), K. *Temnoscheila caerulea* (Olivier, 1790) (photos by A. Samih and N. Maatouf).

Onthophagus maki (Illiger, 1803) (Fig. 5F) [Rc]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 4 individuals, pitfall trap, 01.VI.2022, 4 individuals, pitfall trap, 22.VI.2022, 2 individuals, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution Spain, France, Portugal, Italy, North Africa (Löbl *et al.*, 2006a) and Morocco (Habbaz *et al.*, 2025); Ecological insight: as a dung beetle, *Onthophagus* contributes to nutrient cycling by breaking down dung into smaller pieces, enhancing nutrient availability for plants. The process of dung burial helps incorporate organic matter into the soil (Ramli *et al.*, 2019).

Oxythyrea funesta (Poda, 1761) [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps, 01.VI.2022, 1 individual, pitfall trap, 01.VI.2022, 1 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 2 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps, 21.VII.2022, 26 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, North Africa (Smetana, 2006) and Morocco (Habbaz *et al.*, 2025).

Pleurophorus caesus (Creutzer, 1796)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 3 individuals, pan traps, 12.V.2022, 2 individuals, pitfall trap, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Dellacasa & Dellacasa, 2006).

Protaetia opaca (Fabricius, 1787) [R]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VII.2021, 1 individual, pan traps, 01.VI.2022, 5 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: France, Italy, Portugal, Spain, Canary Islands, Algeria, Tunisia and Morocco (Smetana, 2006).

Scarabaeus cicatricosus (P.H.Lucas, 1846) (Fig. 5C) [Rc]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 21 individuals, pitfall trap, 04.VIII.2021, 4 individuals, pitfall trap, 15.IX.2021, 14 individuals, pitfall trap, 28.X.2021, 12 individuals, pitfall trap, 12.V.2022, 3 individuals, pitfall trap, 01.VI.2022, 4 individuals, pitfall trap, 22.VI.2022, 27 individuals, pitfall trap, 21.VII.2022, 5 individuals, pitfall trap, 30.XII.2022, 1 individual, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 04.VIII.2021, 1 individual, pitfall trap, 28.X.2021, 2 individuals, pitfall trap, 30.XII.2021, 3 individuals, pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, pitfall trap; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Spain, Portugal, Algeria and Morocco (Löbl *et al.*, 2006b; Habbaz *et al.*, 2024); Ecological insight: *Scarabaeus* as a coprophagous species, it feeds primarily on the dung of herbivores, which it collects, buries and consumes (Barbero *et al.*, 1999; Bai *et al.*, 2015). This behavior not only helps to break down and decompose animal waste but also contributes to the redistribution of nutrients back into the soil, promoting soil fertility (Thotagamuwa *et al.*, 2023).

Scarabaeus sacer (Linnaeus, 1758) (Fig. 5B)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 2 individuals, pitfall trap, 15.IX.2021, 1 individual, pitfall trap, 01.VI.2022, 7 individuals, pitfall trap, 21.VII.2022, 1 individual pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, North Africa (Löbl *et al.*, 2006) and Morocco (Habbaz *et al.*, 2025).

Trichius zonatus (Germar, 1831) (Fig. 5G) [R]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 2 individuals, pan traps; Collector name: Samih; Worldwide distribution: Algeria, Tunisia and Morocco (Smetana, 2006).

Tropinota squalida pilosa (Brullé, 1832)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.V.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Asia, North Africa (Smetana, 2006) and Morocco (Habbaz *et al.*, 2025).

Family: Scolytidae Latreille, 1804

Hypoborus ficus (Erichson, 1836)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 2 individuals, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia, North Africa (Knizek, 2011) and Morocco (Habbaz *et al.*, 2025).

Platypus cylindrus (Fabricius, 1792)

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 30.XII.2022, 1 individual, pan traps.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 27.V.2021, 2 individuals, pan traps, 01.VII.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, North Africa (Knizek, 2011) and Morocco (Habbaz *et al.*, 2025).

Xyleborus monographus (Fabricius, 1792) [S]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 04.VIII.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 4 individuals, pan traps, 28.X.2021, 2 individuals, pan traps, 27.IX.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Algeria and Morocco (Knizek, 2011; Habbaz *et al.*, 2025).

Family: Scraptiidae Mulsant, 1856

Scraptia fuscula (Müller, 1821)

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 3 individuals, pan traps, 01.VI.2022, 2 individuals, pan traps, 22.VI.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Algeria and Morocco (Leblanc *et al.*, 2008).

Family: Staphylinidae Latreille, 1802

Oxytelus sculptus (Gravenhorst, 1806) [S]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 2 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, pan traps, 27.IX.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and Morocco (Smetana, 2004).

Philonthus longicornis (Stephens, 1832) [N]

Material examined

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 6 individuals, pan traps, 01.VI.2022, 6 individuals, pan traps, 22.VI.2022, 7 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia and North Africa (Smetana, 2004).

Spedophilus marshami (Stephens, 1832) [N]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 01.VI.2022, 1 individual, pan traps; Collector name: Samih; Worldwide distribution: Europe, Tunisia, South Korea, Russia and Turkey (Smetana, 2004).

Tachyporus hyponorum (Fabricius, 1775)**Material examined**

Zouada, Open forest, 35°10'48.0 N 5°58'40.1 W, 27.V.2021, 1 individual.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 2 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Algeria, Tunisia and Morocco (Smetana, 2004).

Tachyporus nitidulus (Fabricius, 1781) [N]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 2 individuals, pan traps, 12.V.2022, 6 individuals, pan traps, 01.VI.2022, 3 individuals, pan traps; Collector name: Samih; Worldwide distribution: Europe, Asia and North Africa except Morocco (Smetana, 2004).

Xantholinus linearis (Olivier, 1795)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 12.V.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 11 individuals, pan traps, 12.V.2022, 6 individuals, pan traps, 01.VI.2022, 17 individuals, pan traps, 21.VII.2022, 13 individuals, pan traps; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Europe, Asia, North Africa (Smetana, 2004) and Morocco (Habbaz *et al.*, 2025).

Family: Tenebrionidae Latreille, 1802***Adelostoma sulcatum*** (Duponchel, 1829)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 27.IX.202, 1 individual, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Spain, Greece, Asia, North Africa (Löbl *et al.*, 2008) and (Habbaz *et al.*, 2025).

Boromorphus tagenoides (Lucas, 1846)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pitfall trap; Collector name: Maatouf; Worldwide distribution: Spain and North Africa (Löbl *et al.*, 2008).

Cnemeplatia atropos (Costa, 1847)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pitfall trap, 22.VI.2022, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Asia, Tunisia, Egypt and Morocco (Löbl *et al.*, 2008).

Cossyphus hoffmanseggii (Herbst, 1797) (Fig. 5J)**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 04.VIII.2021, 3 individuals, pitfall trap, 15.IX.2021, 1 individual, pitfall trap, 12.V.2022, 1 individual, pitfall trap, 22.VI.2022, 1 individual, pitfall trap, 01.VII.2022, 3 individuals, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution Spain, Portugal, Algeria and Morocco (Löbl *et al.*, 2008).

Gonocephalum granulatum granulatum (Fabricius, 1792) [E]**Material examined**

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pitfall trap; Collector name: Samih; Worldwide distribution: Morocco (Iwan & Löbl, 2008).

Heliotaurus ruficollis tangerianus (Escalera, 1922) [E]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 10 individuals, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 1 individual, pan traps, 12.V.2022, 2 individuals, pan traps, 12.V.2022, 7 individuals, malaise trap, 01.VI.2022, 448 individuals, pan traps, 22.VI.2022, 7 individuals, pan traps, 01.VII.2022, 6 individuals, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Morocco (Novak & Pandtersson, 2008; Habbaz *et al.*, 2025).

Isomira melanophthalma (Lucas, 1846)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 1 individual, imalaise trap; Collector name: Samih & Maatouf; Worldwide distribution: Portugal, France, Italy, Croatia, Bosnia, Algeria, and Morocco (Novak & Pandtersson, 2008; Habbaz *et al.*, 2025).

Latheticus oryzae (Waterhouse, 1880)**Material examined**

Forest edge, 35°12'43.1 N 6°01'57.8 W, 27.V.2021, 1 individual, pan traps; Collector name: Maatouf; Worldwide distribution: Europe, Asia, North Africa (Löbl *et al.*, 2008) and Morocco (Habbaz *et al.*, 2025).

Pimelia chrysomeloides subris (Koch, 1941) [E] [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 15.IX.2021, 13 individuals, pitfall trap, 28.X.2021, 10 individuals, pitfall trap, 12.V.2022, 15 individuals, pitfall trap, 01.VI.2022, 23 individuals, pitfall trap, 22.VI.2022, 8 individuals, pitfall trap, 21.VII.2022, 11 individuals, pitfall trap, 27.IX.2022, 11 individuals, pitfall trap, 30.XII.2022, 9 individuals, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 2 individuals, pitfall trap, 15.IX.2021, 4 individuals, pitfall trap, 28.10.2021, 3 individuals, pitfall trap, 30.XII.2021, 4 individuals, pitfall trap; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Morocco (Löbl *et al.*, 2008; Habbaz *et al.*, 2025); Ecological insight: Genus *Pimelia* is decomposer (saprophagous), primarily feeding on decaying plant material, seeds and occasionally fungi (De Vega, 2011; Karmaoui *et al.*, 2023).

Sepidium bidentatum (Solier, 1843)**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VI.2022, 01.VII.2021, 4 individuals, pitfall trap, 12.V.2022, 6 individuals, pitfall trap, 01.VI.2022, 27 individuals, pitfall trap, 22.VI.2022, 4 individuals, pitfall trap.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 12.V.2022, 4 individuals, pitfall trap, 22.VI.2022, 1 individual, pitfall trap; Collector name: Samih & Maatouf; Worldwide distribution: Spain, Portugal and Morocco (Löbl *et al.*, 2008).

Zophosis minuta (Fabricius, 1775) [S]**Material examined**

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 01.VII.2021, 12 individuals, pitfall trap, 04.VIII.2021, 1 individual, pitfall trap, 15.IX.2021, 7 individuals, pitfall trap, 28.X.2021, 4 individuals, pitfall trap, 12.V.2022,

24 individuals, pitfall trap, 01.VI.2022, 9 individuals, pitfall trap, 22.VI.2022, 7 individuals, pitfall trap, 21.VII.2022, 3 individuals, pitfall trap, 27.IX.2022, 2 individuals, pitfall trap.

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 01.VII.2021, 1 individual, pitfall trap, 04.VIII.2021, 4 individuals, pitfall trap, 28.X.2021, 1 individual, pitfall trap; Collector name: Samih, Maatouf & Habbaz; Worldwide distribution: Tunisia (Lefèvre, 1885), Portugal, Spain, Morocco (Löbl *et al.*, 2008; Habbaz *et al.*, 2025).

Family: Trogossitidae Latreille, 1802

Temnoscheila caerulea (Olivier, 1790) (Fig. 5K) [N] [R]

Material examined

Larache, Dense forest, 35°11'40.9 N 06°07'34.9 W, 15.IX.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Canary Islands, Tunisia and China (Kolibac, 2007); Ecological insight: An important predator of bark beetles (Cebeci & Baydemir, 2018). It helps regulate their populations, reducing the risk of infestations and preventing potential damage to trees and forest ecosystems (Ettxebeste *et al.*, 2013).

Family: Zopheridae Solier, 1834

Endophloeus markovichianus (Piller & Mitterpacher, 1783) (Fig. 5H) [R]

Material examined

Zouada, Open forest, 35°03'20.2 N 6°01'48.0 W, 15.IX.2021, 1 individual, pan traps.

Rissana, Forest edge, 35°12'43.1 N 6°01'57.8 W, 15.IX.2021, 1 individual, pan traps; Collector name: Samih & Maatouf; Worldwide distribution: Europe, Algeria, Tunisia and Morocco (Ślipinski & Schuh, 2008).

Discussion

Beetle diversity and taxonomic insights

The results aim to address the gap in scientific knowledge regarding beetle communities in Moroccan forest ecosystems, particularly within the cork oak forest of Larache, located on the Atlantic coastal plain. A total of 180 have been identified at the species level, belonging to 37 families, and have been inventoried. It can be deduced that there is significant taxonomic diversity among beetles, compared to increasingly anthropized environments. In the northeastern cork oak forests of Algeria, beetles are represented by 74 species (Ganaoui *et al.*, 2020). In contrast, 216 species belonging to 42 families have been recorded in the cork oak forest of Maâmora (Habbaz *et al.*, 2025). This difference could be partly explained by the greater fragmentation of our study area compared to Maâmora, which may influence the diversity and distribution of species. Additionally, it is important to note that the use of Malaise traps in our study, while effective for certain groups, may not capture as many beetle species as interception traps. The latter were used in the study of Maâmora (Habbaz *et al.*, 2025) and are particularly effective for capturing beetle species, especially saproxylic groups, which are associated with decaying wood and contribute significantly to overall diversity.

Ecological significance of saproxylic and red list species

In our study, we inventoried only 13 saproxylic species, which confirms, on one hand, that the Larache cork oak forest is an ecosystem undergoing changes with unprotected habitats, and on the other hand, it is possible our study sites did not have enough dead wood or old trees to support robust saproxylic insect populations. However, the limited occurrence of saproxylic species may be attributed to the sampling methods employed, as these species are commonly collected using flight interception traps (Bouget *et al.*, 2008; Burner *et al.*, 2022b), techniques not employed in this study. Ten species of beetles are now listed in the Mediterranean Red List, highlighting their status as rare and threatened species. Within this group, 7 saproxylic species are notable for being listed in the Red List of Saproxylic Beetles of the Mediterranean region, while 3 are listed as coprophagous in the Red List of coprophagous species in the Mediterranean region. Saproxylic species, which depend on dead or decaying wood at some stage of their life cycle, play a vital ecological role in forest ecosystems (Seibold & Thorn, 2018; Zúmr *et al.*, 2024).

Table 2. List of species collected in the Cork Oak forest of Larache at sites Dense forest (S1), Open Forest (S2) and Forest edge (S3), and their status: [S] Saproxyl species, [R] Saproxyl species and [Rc] coprophagous species; listed in the Red List of the Mediterranean region, [N] Species newly recorded for Morocco and [E] Endemic species.

Families	Species	Status	Dense forest	Forest edge	Open Forest
Anthicidae	<i>Hirticollis quadriguttatus</i> (Rossi, 1792)			*	
	<i>Omonadus bifasciatus</i> (Rossi, 1792)			*	
	<i>Omonadus floralis</i> (Linnaeus, 1758)			*	
Brentidae	<i>Apion frumentarium</i> (Walton, 1844)	N	*	*	
	<i>Hemitrichapion reflexum</i> (Gyllenhal, 1833)	N	*	*	
Bruchidae	<i>Bruchidius biguttatus</i> (Olivier, 1795)		*		
	<i>Bruchidius bimaculatus</i> (Olivier, 1795)			*	
	<i>Bruchidius foveolatus</i> (Gyllenhal, 1833)		*	*	
Buprestidae	<i>Acmaeodera adasperula</i> (Illiger, 1803)	S	*	*	
	<i>Acmaeodera discoidea</i> (Fabricius, 1787)			*	
	<i>Acmaeodera lanuginosa</i> (Gyllenhal, 1817)		*	*	
	<i>Agrilus biguttatus</i> (Fabricius, 1777)	S		*	
	<i>Agrilus graminis mamorensis</i> (Théry, 1930)	E, S			*
	<i>Anthaxia millefolii polychloros</i> (Abeille de Perrin, 1894)		*	*	
	<i>Anthaxia scutellaris atlasica</i> (Théry, 1930)				*
	<i>Anthaxia umbellatarum umbellatarum</i> (Fabricius, 1787)		*	*	
	<i>Habroloma triangulare</i> (Lacordaire, 1835)		*		
Carabidae	<i>Acupalpus brunnipes</i> (Sturm, 1825)			*	
	<i>Agonum emarginatum</i> (Gyllenhal, 1827)	N		*	
	<i>Bembidion biguttatum</i> (Fabricius, 1779)	N		*	
	<i>Bembidion bipunctatum laevifrons</i> (Schaufuss, 1882)			*	
	<i>Bradycellus verbasci</i> (Duftschmid, 1812)			*	
	<i>Carterus rotundicollis</i> (Rambur, 1837)			*	
Carabidae	<i>Cincidela Moroccana</i> (Fabricius, 1801)		*		
	<i>Cymindis lineola</i> (L. Dufour, 1820)	N	*		
	<i>Cymindis platycollis</i> (Say, 1823)	N			*
	<i>Harpalus attenuatus</i> (Stephens, 1828)			*	
	<i>Microlestes abeillei brisouti</i> (Holdhaus, 1912)			*	
	<i>Ophonus ardosiacus</i> (Lutshnik, 1922)		*		
	<i>Ophonus subquadratus</i> (Dejean, 1829)	N		*	
	<i>Paradromius linearis</i> (Olivier, 1795)			*	
	<i>Philorhizus notatus</i> (Stephens, 1827)	N	*		
	<i>Pterostichus elongatus</i> (Duftschmid, 1812)		*	*	*
	<i>Scarites terricola terricola</i> (Bonelli, 1813)		*		
	<i>Singilis soror soror</i> (Rambur, 1837)		*		
	<i>Stenolophus abdominalis abdominalis</i> (Gene, 1836)		*		
	<i>Stenolophus teutonius</i> (Schränk, 1781)			*	
	<i>Syntomus foveatus</i> (Geoffroy, 1785)			*	
Cerambycidae	<i>Agapanthia irrorata</i> (Fabricius, 1787)			*	
	<i>Alocerus moesiacus</i> (Frivaldszky, 1837)				*
	<i>Certallum ebulinum</i> (Linné, 1767)			*	
	<i>Chlorophorus fávieri</i> (Farimaire, 1873)	E, R		*	
	<i>Phytoecia coerulescens</i> (Scopoli, 1763)			*	
	<i>Oxypleurus nodieri</i> (Mulsant, 1839)			*	
	<i>Stenurella approximans</i> (Rosenhauer, 1856)		*	*	
	<i>Stictoleptura fontenayi</i> (Mulsant and Rey, 1839)		*		
	<i>Trichoferus ilicis</i> (Sama, 1987)	E, R	*	*	
Chrysomelidae	<i>Aphthona euphorbiae</i> (Schränk, 1781)			*	

Table 2. Countinued

	<i>Cassida vittata</i> (Villers, 1789)		*	
	<i>Chrysolina bankii</i> (Fabricius, 1775)		*	
	<i>Chrysolina diluta</i> (Germar, 1823)	N		*
	<i>Cryptocephalus fulvus</i> (Goeze, 1777)	N	*	*
	<i>Cryptocephalus numidicus</i> (Bourdonné, 1876)		*	
	<i>Longitarsus aeneus</i> (Kutschera, 1862)		*	*
	<i>Longitarsus ochroleucus</i> (Marsham, 1802)		*	*
	<i>Oulema melanopus</i> (Linnaeus, 1758)			*
	<i>Psylliodes cuprea</i> (Koch, 1803)		*	
Coccinellidae	<i>Adalia decempunctata</i> (Linnaeus, 1758)			*
	<i>Chilocorus bipustulatus</i> (Linnaeus, 1758)			*
	<i>Coccinella septempunctata</i> (Linnaeus, 1758)			*
	<i>Oenopia conglobata</i> (Linnaeus, 1758)		*	*
	<i>Oenopia lyncea lyncea</i> (Olivier, 1808)		*	
	<i>Platynaspis luteorubra</i> (Goeze, 1777)		*	
	<i>Rhyzobius litura</i> (Fabricius, 1787)		*	*
	<i>Rhyzobius lophantae</i> (Blaisdell, 1892)		*	*
	<i>Rodolia cardinalis</i> (Mulsant, 1850)		*	*
	<i>Scymnus abiandis</i> (Paykull, 1798)	N		*
	<i>Scymnus apandzi</i> (Mulsant, 1846)	N	*	*
	<i>Scymnus impexus</i> (Mulsant, 1850)	N		*
	<i>Scymnus interruptus</i> (Goeze, 1777)		*	*
	<i>Scymnus subvillosus</i> (Goeze, 1777)			*
	<i>Scymnus suturalis</i> (Westman, 1795)	N	*	
	<i>Standhorus punctillum</i> (Weise, 1891)		*	
Corylophidae	<i>Arthrolips convexiuscula</i> (Motschulsky, 1849)			*
Curculionidae	<i>Brachyderes incanus</i> (Linné, 1758)		*	*
	<i>Brachyderes pubescens</i> (Boheman, 1833)		*	*
	<i>Brachytemnus porcatus</i> (Germar, 1823)		*	*
	<i>Ceutorhynchus pallidactylus</i> (Marsham, 1802)		*	
	<i>Coeliodes ruber</i> (T.Marsham, 1802)		*	
	<i>Lixus juncii</i> (Boheman, 1835)	N	*	*
	<i>Microplontus rugulosus</i> (J.F.W.Herbst, 1795)		*	
	<i>Orchestes irroratus maroccanus</i> (Roudier, 1954)		*	
Curculionidae	<i>Sitona callosus</i> (Gyllenhal, 1834)	N		*
	<i>Sitona lineatus</i> (Linnaeus, 1758)	N	*	
	<i>Sitona lineellus</i> (Bonsdorff, 1785)	N	*	
	<i>Sitona longulus</i> (Gyllenhal, 1834)	N	*	*
	<i>Tychius cuprifer</i> (Panzer, 1799)			*
	<i>Tychius pusillus</i> (Germar, 1842)		*	
Dasytidae	<i>Dasytes nigroaeneus</i> (Küster, 1850)		*	*
	<i>Dasytes terminalis</i> (Jacquelin du Val, 1863)		*	*
Dermestidae	<i>Anthrenus flavipes</i> (LeConte, 1854)		*	
	<i>Anthrenus fuscus</i> (Olivier, 1790)		*	*
	<i>Anthrenus museorum</i> (Linnaeus, 1761)		*	
	<i>Anthrenus pimpinellae</i> (Fabricius, 1775)			*
	<i>Attagenus bifasciatus</i> (Fabricius, 1787)		*	*
	<i>Dermestes frischii</i> (Kugelann, 1792)		*	*
	<i>Orphilus niger</i> (Rossi, 1790)		*	
	<i>Trogoderma granarium</i> (Everts, 1898)		*	*
Dryopidae	<i>Dryops luridus</i> (Erichson, 1847)	N	*	
Elateridae	<i>Cardiophorus rufipes</i> (Buysson, 1902)	S	*	*
	<i>Drasterius bimaculatus</i> (Rossi, 1790)	S	*	
Erotylidae	<i>Triplax lacordairei</i> (Crotch, 1870)	R	*	
Geotrupidae	<i>Thorectes distinctus</i> (Marseul, 1878)		*	*
	<i>Typhaeus typhoeus</i> (Linnaeus, 1758)			*
Glaphyridae	<i>Anthypna meles</i> (Fabricius, 1792)		*	*

Table 2. Countinued.

	<i>Eulasia goudoti</i> (Laporte, 1840)	E	*		
Histeridae	<i>Hypocaccus rugiceps</i> (Duftschmid, 1805)			*	
	<i>Saprinus proximus similimus</i> (Wollaston, 1865)		*		
Hydraenidae	<i>Ochthebius bicolon</i> (Germar, 1823)	N			*
Hydrophilidae	<i>Cercyon obsolendus</i> (Gyllenhal, 1808)	N		*	
	<i>Sphaeridium scarabaeoides</i> (Linnaeus, 1758)	N		*	
Laemophloeidae	<i>Laemophloeus monilis</i> (Fabricius, 1787)	S			*
	<i>Laemophloeus muticus</i> (Fabricius, 1781)	N		*	
Latridiidae	<i>Corticarina curta</i> (Wollaston, 1854)			*	
	<i>Enicmus transversus</i> (Olivier, 1790)		*		
	<i>Stephostethus productus</i> (Rosenhauer, 1856)			*	
Meloidae	<i>Berberomeloe majalis</i> (Linnaeus, 1758)		*		
	<i>Croscherichia paykulli</i> (Billberg, 1813)		*		
	<i>Hycleus rufipalpis</i> (Escalera, 1909)		*		
	<i>Mylabris variabilis</i> (Pallas, 1781)			*	
Melolonthidae	<i>Euserica mamorensis</i> (Baraud, 1965)	E	*	*	*
	<i>Hoplia africana</i> (Escalera, 1914)			*	
	<i>Hoplia bilineata</i> (Fabricius, 1801)			*	
	<i>Hoplia philanthus</i> (Fuessly, 1775)		*	*	
	<i>Sphodroxia Moroccana</i> (Ley, 1923)	E			
Melyridae	<i>Aplocnemus virens</i> (Suffrian, 1843)	N	*	*	
	<i>Colotes javeti</i> (Jacquelin du Val, 1852)		*		
	<i>Psilothrix viridicoerulea</i> (Geoffroy, 1758)		*	*	
	<i>Troglops furcatus</i> (Perrin, 1885)			*	*
Mordellidae	<i>Mediimorda bipunctata</i> (Germar, 1827)			*	*
	<i>Mordella aculeata</i> (Linnaeus, 1758)		*	*	
	<i>Variimorda villosa</i> (Schränk von Paula, 1781)	N	*	*	
Nitidulidae	<i>Carpophilus hemipterus</i> (Linnaeus, 1758)			*	*
	<i>Eपुरaea latipes</i> (Grouvelle, 1896)			*	
	<i>Meligethes aeneus</i> (Fabricius, 1775)		*	*	*
	<i>Meligethes viridescens</i> (Fabricius, 1787)		*	*	*
Oedemeridae	<i>Chrysanthia viridissima</i> (Linnaeus, 1758)	N	*		
	<i>Oedemera barbara</i> (Fabricius, 1792)	S	*	*	*
	<i>Oedemera marmorata</i> (Erichson, 1841)			*	
Phalacridae	<i>Olibrus pygmaeus</i> (Sturm, 1807)		*	*	*
	<i>Phalacrus coruscus</i> (Panzer, 1797)			*	
Ptinidae	<i>Dignomus irroratus</i> (Kiesenwandter, 1851)	N	*	*	
	<i>Stegobium paniceum</i> (Linnaeus, 1758)	S	*	*	
Scarabaeidae	<i>Anisoplia baetica</i> (Erichson, 1847)			*	
	<i>Anthoplia floricola</i> (Fabricius 1787)			*	
	<i>Aphodius diecki</i> (Harold, 1870)			*	
	<i>Blitopertha lineata</i> (Fabricius, 1798)			*	
	<i>Gymnopleurus flagellatus</i> (Fabricius, 1787)		*		
	<i>Gymnopleurus sturmii</i> (MacLeay, 1821)	Rc	*		
	<i>Onthophagus maki</i> (Illiger, 1803)	Rc	*		
	<i>Oxythyrea funesta</i> (Poda, 1761)	S	*	*	
	<i>Pleurophorus caesus</i> (Creutzer, 1796)			*	
	<i>Protaetia opaca</i> (Fabricius, 1787)	R	*	*	
	<i>Scarabaeus cicatricosus</i> (P.H.Lucas, 1846)	Rc	*	*	*
	<i>Scarabaeus sacer</i> (Linnaeus, 1758)		*		
	<i>Trichius zonatus</i> (Germar, 1831)	R		*	
	<i>Tropinota squalida pilosa</i> (Brullé, 1832)		*	*	
Scolytidae	<i>Hypoborus ficus</i> (Erichson, 1836)			*	
	<i>Platypus cylindrus</i> (Fabricius, 1792)		*		*
	<i>Xyleborus monographus</i> (Fabricius, 1792)	S	*	*	
Scraptiidae	<i>Scraptia fuscula</i> (Müller, 1821)			*	
Staphylinidae	<i>Oxytelus sculptus</i> (Gravenhorst, 1806)	S	*	*	

Table 2. countinued

	<i>Philonthus longicornis</i> (Stephens, 1832)	N	*	
	<i>Spedophilus marshami</i> (Stephens, 1832)	N	*	
	<i>Tachyporus hyponorum</i> (Fabricius, 1775)		*	*
	<i>Tachyporus nitidulus</i> (Fabricius, 1781)	N	*	
	<i>Xantholinus linearis</i> (Olivier, 1795)		*	*
Tenebrionidae	<i>Adelostoma sulcatum</i> (Duponchel, 1829)		*	*
	<i>Boromorphus tagenoides</i> (Lucas, 1846)		*	
	<i>Cnemeplatia atropos</i> (Costa, 1847)		*	
	<i>Cossyphus hoffmanseggii</i> (Herbst, 1797)		*	
	<i>Gonocephalum granulatum granulatum</i> (Fabricius, 1792)	E	*	
	<i>Heliotaurus ruficollis tangerianus</i> (Escalera, 1922)	E	*	*
	<i>Isomira melanophthalma</i> (Lucas, 1846)		*	*
	<i>Latheticus oryzae</i> (Waterhouse, 1880)			
	<i>Pimelia chrysomeloides subris</i> (Koch, 1941)	E, S	*	*
	<i>Sepidium bidentatum</i> (Solier, 1843)		*	*
	<i>Zophosis minuta</i> (Fabricius, 1775)	S	*	*
Trogossitidae	<i>Temnoscheila caerulea</i> (Olivier, 1790)	N, R	*	*
Zopheridae	<i>Endoploeus markovichianus</i> (Piller & Mitterpacher, 1783)	R	*	*

They contribute to nutrient cycling and organic matter decomposition, facilitating soil formation and forest habitat regeneration (Parisi *et al.*, 2020). These species are also recognized as valuable bioindicators of forest health, as their presence reflects the availability of deadwood and the continuity of forest ecosystems (Tsikas & Karanikola, 2020). Cavities and crevices in old trees, along with deadwood, provide essential breeding sites and food sources for these organisms (Micó, 2018). The rarity and threatened status of species provide valuable insights into the ecological balance and uniqueness of the cork oak forest in Larache amidst various anthropogenic pressures. These rare taxa are highly sensitive to environmental disturbances, with their often restricted distribution within geographic ranges making them ecologically valuable. They occupy specific habitats and have particular trophic requirements (Caprenato, 2015). The loss of this sensitive group can have negative impacts on ecosystem functioning, emphasizing the importance of conservation measures to ensure the sustainability of this ecosystem (Burner *et al.*, 2022a).

Endemic and new species

During this study, we identified 9 endemic species with limited spatial distribution capacities. These endemic species provide a unique perspective on the complexity and specificity of the cork oak forest ecosystem in Larache. Although their distribution may be restricted to specific areas, their local abundance underscores their key role in preserving the biological diversity of the region (Iorio *et al.*, 2022). These endemic species play a crucial role in maintaining biodiversity by contributing to the richness and complexity of the local flora and fauna, which are essential for sustaining a balanced and diverse ecological community. They often occupy specific ecological niches, interacting with other species such as pollinators, decomposers and predators, thereby helping to preserve the ecological functions of the cork oak forest of Larache.

Our inventory also reveals 33 species new to Morocco, according to the catalogue of beetle distribution in the Palaearctic region. It is important to clarify that these species are not necessarily "new" in the strict sense, but rather new to the Moroccan and regional context due to the lack of detailed prior inventories. This situation can be explained by several factors. On one hand, the region has undergone significant anthropogenic transformations since the 12th century, such as changes in land use, urbanization, and climate change (Ballouche, 2013), which have altered natural habitats and facilitated the appearance or migration of new species. On the other hand, some species may have been present in the region for a long time but were not previously identified due to a lack of reliable historical data or comprehensive surveys. Additionally, natural species redistributions in response to environmental changes could also explain the increased presence of certain species.

Conservation implications and recommendations

The heritage value of the cork oak forest in Larache underscores the necessity to implement effective protection measures against threats such as overgrazing, agricultural clearing, and excessive collection of deadwood. It is essential to enhance our understanding of the biology and behavior of the identified species. The preservation and restoration of suitable habitats for active beetle species are crucial. This entails protecting forested areas, ensuring an adequate amount of deadwood is left in situ and avoiding excessive clear-cutting, which supports the creation of complex and diverse microhabitats for beetles. Conservation management of forest ecosystems as a whole requires the designation of isolated, low-disturbance areas as biodiversity refuges (Gosselin & Pailland, 2017). These measures should be developed in close collaboration with management authorities, local communities, and stakeholders to ensure an effective approach for long-term biodiversity conservation across forested areas. It is therefore essential to establish a specific Red List of threatened species in Morocco and to periodically revise it. Creating such a list would significantly contribute to the conservation of regional biodiversity and guide management policies towards targeted actions. As key community decision-makers, their support for establishing a Red List for beetles enhances recognition of the heritage value of the Larache cork oak forest and demonstrates their commitment to preserving this environmental legacy.

Author's Contributions

Amine Samih: methodology, formal analysis, investigation, draft preparation, final review and edit, visualization and supervision. **Noureddin Maatouf:** supervision, methodology, project administration and funding acquisition. **Sergi Trócoli:** confirmed identification, final review and edit. **Hamza Habbaz:** formal analysis and references data. **Latifa Rohi:** supervision, project administration, final review and edit.

Author's Information

Amine Samih	✉ aminesamih96@gmail.com	 https://orcid.org/0009-0003-0480-7119
Noureddin Maatouf	✉ noureddin_maatouf@yahoo.ca	 https://orcid.org/0000-0002-6288-7378
Sergi Trócoli	✉ sergitrocoli@gmail.com	 https://orcid.org/0009-0005-2643-096X
Hamza Habbaz	✉ Hamza96habbaz@gmail.com	 https://orcid.org/0009-0000-7117-1075
Latifa Rohi	✉ rohilatifa@gmail.com	 https://orcid.org/0000-0002-4180-1117

Funding

This research received no specific grant from any funding agencies.

Data Availability Statement

The specimens collected was deposited in the insect collection of the Centre for Innovation, Research, and Training - National Agency for Water and Forests, BP. 763, Rabat - Agdal, Morocco.

Acknowledgments

We extend our heartfelt thanks to the Director of the Centre for Innovation, Research, and Training in Rabat (National Agency for Water and Forests) for their hospitality and unwavering support in facilitating this work. We are also grateful to El Ayyachi El Mnouar (Technician in the field of forest management - National Agency for Water and Forests Rabat, Morocco) for their guidance and assistance. Special thanks to Dr. Piero Leo (Private entomologist, Cagliari, Italy) for his invaluable identification of some specimens. We thank also the reviewers and editors for their constructive comments.

Ethics Approval

Beetles were used in this study. All applicable international, national, and institutional guidelines for the care and use of animals were followed. This article does not contain any studies with human participants performed by any of the authors.

Conflict of Interest

The author declares that there is no conflict of interest regarding the publication of this paper.

REFERENCES

- Alizadeh, M., Safaie, N., Azarbad, H., Mehrabadi, M. & Shamsbakhsh, M. (2024) Detection of *Paecilomyces formosus* associated with declining urban forests and beetles in Iran. *Physiological and Molecular Plant Pathology*, 133, 102384. <https://doi.org/10.1016/j.pmpp.2024.102384>
- Alonso, C. (2007) Coleopteres Chrysomelidae Clytrinae de France continentale et de corse. <http://www.insecte.org/archives/CLYTRINAE>
- Amokrane, A., Khammar, H., Hadjab, R. & Saheb, M. (2020) Evaluation of the Diversity of Durum Wheat Coleoptera (*Triticum Durum* Desf.) in the Region of Sigus Oum El Bouaghi (Eastern Algeria). *Journal of Bioresource Management*, 7(4), 33–54. <https://doi.org/10.35691/jbm.0202.0149>
- Anton, K. W. (2010) Bruchinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 6. Apollo Books, Stenstrup, 339353.
- Arahou, M. (2008) Catalogue de l'entomofaune du Chêne vert du Moyen Atlas (Maroc). Rabat, Maroc, Institut Scientifique, N°22, 9. ISBN: 9954-8347-8-8.
- Arce-Pérez, R., Gómez-Anaya, J. A. & Epler, J. H. (2018) The Family Noteridae Thomson (Coleoptera: Adephaga) in Miradores Lagoon, Veracruz, Mexico, with a Description of its Assemblage. *The Coleopterists Bulletin*, 72, 75-83. <https://doi.org/10.1649/0010-065X-72.1.75>
- Bai, M., Li, S., Lu, Y., Yang, H., Tong, Y. & Yang, X. (2015) Mandible evolution in the Scarabaeinae (Coleoptera: Scarabaeidae) and adaptations to coprophagous habits. *Frontiers in Zoology*, 12, 1-10. <https://doi.org/10.1186/s12983-015-0123-z>
- Bajocco, S., De Angelis, A., Perini, L., Ferrara, A. & Salvati, L. (2012) The impact of land use/land cover changes on land degradation dynamics: a Mediterranean case study. *Environmental Management*, 49, 980-989. <https://doi.org/10.1007/s00267-012-9831-8>
- Balkenohl, M. (2003) Carabidae, Tribe: Scaritini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 231234.
- Ballouche, A. (2013) Contribution à l'histoire récente de la végétation du Bas-Loukkos (province de Larache, Morocco). *Physio-Géo*, 7, 1-452. <https://doi.org/10.4000/physio-geo.3426>
- Ballouche, A. (2024) Missing Landscapes: A Geohistory of Parkland Landscapes in Northwestern Morocco. *Land*, 13(5), 649. <https://doi.org/10.3390/land13050649>
- Baraud, J. (1992) Encart Coléoptères Scarabaeoidea d'Europe. *Publications de la Société Linnéenne de Lyon*, 61(9), 1-2. <https://doi.org/10.3406/linly.1992.11007>
- Barbero, E., Palestini, C. & Rolando, A. (1999) Dung beetle conservation: effects of habitat and resource selection (Coleoptera: Scarabaeoidea). *Journal of Insect Conservation*, 3, 75-84. <https://doi.org/10.1023/a:1009609826831>
- Bellifa, M. & Roussand, J. (2021) Sur la présence en France de *Dasytes terminalis* Jacquelin du Val, 1863 avec des données inédites sur sa répartition and un point sur sa biologie, (Coleoptera, Dasytidae). *Le Coléoptériste*, 24(2), 128–131. <https://hal.science/hal-03466070>
- Benabdelkader, M., Saifi, R. & Saifi, H. (2021) Sustainable Agriculture in Some Arab Maghreb Countries (Morocco, Algeria, Tunisia). *Agro-Environmental Sustainability in MENA Regions*, 233-261. https://doi.org/10.1007/978-3-030-78574-1_10
- Benyahia, Y., Latifa, R., Valladarès, L., Maatouf, N., Courtin, O., El Antry, S. & Brustel, H. (2016) Preliminary list of Coleoptera heritage species of the Talassemtane National Park, Morocco. *Journal of Insect Biodiversity*, 4(13), 1-30. <https://doi.org/10.12976/jib/2015.4.13>
- Bezdek, A. (2006) Melolonthidae: Tribe: Melolonthini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 191-198.
- Bily, S. (2006) Buprestidae, Buprestinae: Trib. Anthaxiini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 369-381.
- Bologna, M. A. (2008) Meloidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 370-412.
- Borowiec, L. & Sekerka, L. (2010) Cassidinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 6. Apollo Books, Stenstrup, 368-390.

- Borowski, J. (2007) Ptinidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 328-362.
- Bouget, C., Brustel, H., Brin, A. & Noblecourt, T. (2008) Sampling saproxylic beetles with window flight traps: methodological insights. *Revue d'Ecologie (La Terre et la Vie)*, 21-32. <https://doi.org/10.3406/revec.2008.1457>
- Bousquet, Y. (2003a) Carabidae: Tribe Platynini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 449-454.
- Bousquet, Y. (2003b) Carabidae: Tribe Pterostichini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 469-515.
- Bowstead, S., (2007). Corylophidae, pp. 631-635. In: Löbl, I. & Smetana, A. (eds.), Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 935.
- Brendell, M. J. D. (1975) Handbooks for the identification of British insects (Coleoptera Tenebrionidae), 5, 1-22.
- Burner, R. C., Drag, L., Stephan, J. G., Birkemoe, T., Wandherbee, R., Muller, J. & Thygeson, A. (2022a) Functional structure of European forest beandle communities is enhanced by rare species. *Biological Conservation*, 267, 109491. <https://doi.org/10.1016/j.biocon.2022.109491>
- Burner, R. C., Birkemoe, T., Åström, J. & Thygeson, A. (2022b) Flattening the curve: approaching complete sampling for diverse beetle communities. *Insect Conservation and Diversity*, 15(2), 157-167. <https://doi.org/10.1111/icad.12540>
- Caldara, R. (2013) Curculioninae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 8. Leiden, Brill, 117-172. https://doi.org/10.1163/9789004259164_011
- Carpaneto, G. M., Baviera, C., Biscaccianti, A. B., Brandmayr, P., Mazzei, A., Mason, F., Battistoni, A., Teofili, C., Rondinini, C., Fattorini, S. & Audisio, P. A. (2015) A red list of Italian Saproxylic beetles: taxonomic overview, ecological features and conservation issues (Coleoptera). *Fragmenta Entomologica*, 47(2), 53-126. <https://doi.org/10.4081/fe.2015.138>
- Carrillo, J. G., García, P. C., Mendez, J. L. T., Barreda, J. M., Pérez, C. & Bensusan, K. (2018) Los Anthicidae Latreille, 1819 (Insecta, Coleoptera) de la provincia de Cádiz (Andalucía, S. de la Península Ibérica). *Revista Gaditana De Entomología*, 9(1), 121-149.
- Cate, P. C. (2007) Elateridae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 94-209.
- Cálix, M., Alexander, K. N. A., Nieto, A., Dodelin, B., Soldati, F., Telnov, D., VazquezAlbalade, X., Aleksandrowicz, O., Audisio, P., Istrate, P., Jansson, N., Legakis, A., Liberto, A., Makris, C., Merkl, O., Mugerwa Pettersson, R., Schlaghamersky, J., Bologna, M.A., Brustel, H., Buse, J., Novák, V. & Purchart, L. (2018) European Red List of Saproxylic Beetle. Supplementary Material to the IUCN European Red List of Saproxylic Beetles. Brussels, Belgium: IUCN, 1-15. <https://www.iucnredlist.org/resources/calix2018>
- Cebeci, H. H. & Baydemir, M. (2018) Depredadores de escarabajos de la corteza (Coleoptera) en la región de Balıkesir de Turquía. *Revista Colombiana de Entomología*, 44(2), 283-287. <https://doi.org/10.25100/socolen.v44i2.7326>
- Chandler, D. S., Uhmman, G., Nardi, G. & Telnov, D. (2008) Anthicidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 421-455.
- Charrier, S. (2002) Clé de détermination des Coléoptères Lucanides et Scarabéides de Vendée et de l'Ouest de la France. *Le Naturaliste Vendéen*, N°2, 61-93.
- Chavanon G. (2018) Catalogue des Coléoptères de la région Orientale du Maroc (Province de Guercif exceptée). *Travaux de l'Institut Scientifique, Série Zoologie*, 57, 1-193.
- Colonnelli, E. (2013) Centorhynchinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 8. Leiden, Brill, 176-213.
- Davis, A. J., Holloway, J. D., Huijbregts, H., Krikken, J., Kirk-Spriggs, A. H. & Sutton, S. L. (2001) Dung beetles as indicators of change in the forests of northern Borneo. *Journal of Applied Ecology*, 38(3), 593-616. <https://doi.org/10.1046/j.1365-2664.2001.00619.x>
- Delbol, M. & Lempereur, J. M. (2014) Apport à la connaissance des Sisonini de Belgique (Curculionidae: Entiminae). *Entomologie Faunistique*, 67, 15-25.
- Delfosse, E. (2009) Le nombre d'espèces d'Insectes connus en France et dans le monde (Arthropoda: Insectes). *Le Bulletin d'Arthropoda*, 42, 4-37.
- Dellacasa, M. & Dellacasa, G. (2006) Aphodiinae: Tribe Aphodiini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 105-142.
- De Vega, C., Arista, M., Ortiz, P.L., Herrera, C. M. & Talavera, S. (2011) Endozoochory by beetles: a novel seed dispersal mechanism. *Annals of Botany*, 107(4), 629-637. <https://doi.org/10.1093/aob/mcr013>

- Doberl, M. (2010) Alticinae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 6, Apollo Books, Stenstrup, 491-563.
- Dubucq, C. & Jacquemart, A. L. (2020) Influences de la composition florale des biotopes en paysages agricoles sur les interactions plantes-pollinisateurs. Master en sciences agronomiques, 118.
- Ehret, J. M. (1990) Les Apions de France: Clés d'identification commentées (Coleoptera Curculionidae Apioninae). *Bulletin de la Société Linnéenne de Lyon*, 59(7), 209-292. <https://doi.org/10.3406/linly.1990.14241>
- Ekschmitt, K., Weber, M. & Wolters, V. (1997) Spiders, carabids, and staphylinids: the ecological potential of predatory macroarthropods. *Fauna in Soil Ecosystems*, CRC Press, 321-376. <https://doi.org/10.1201/9781482273571-10>
- Etxebeste, I., Lencina, J. L. & Pajares, J. (2013) Saproxylic community, guild and species responses to varying pheromone components of a pine bark beetle. *Bulletin of Entomological Research*, 103(5), 497-510. <https://doi.org/10.1017/S0007485312000879>
- Fadda, S. (2016) Les Oedemeridae Latreille, 1810 de la faune de France: clé de détermination et éléments d'écologie et de biologie (Coleoptera Tenebrionidea). *Naturalia-Environnement*, L'Entomologiste, 72(3), 141-155.
- Farag, M. R., Anter, R. G., Elhady, W. M., Khalil, S. R., Abou-Zeid, S. M. & Hassanen, E. A. (2021) Diversity, succession pattern and colonization of forensic entomofauna on indoor rat carriers concerning the manner of death. *Rendiconti Lincei, Scienze Fisiche e Naturali*, 32, 521-538. <https://doi.org/10.1007/s12210-021-01009-w>
- Fernandes, M. E., Ferreira, C. & Figueiredo, E. (2024) Participatory methods and approaches in the management of Mediterranean forests: a systematic literature review. *International Forestry Review*, 26(2), 177-197. <https://doi.org/10.1505/146554824838819851>
- Ferrer, J., Tovar, A. C. & Gutiérrez, J. L. (2009) Comentarios sobre Cnemeplatini Jacquelin du Val, 1861 (Coleoptera, Tenebrionidae) de la península Ibérica e islas Canarias. *Graellsia*, 65(2), 133-144. <https://doi.org/10.3989/graellsia.2009.v65.i2.146>
- Ganaoui, N. & Chakali, G. (2023) Diversité and importance des Coléoptères dans les forêts de chênes (nord-est de l'Algérie). *IOBC-WPRS Bullandin*, 168, 16-23.
- García, N., Numa, C., Bartolozzi, L., Brustel, H., Buse, J., Norbiato, M., Recalde, J. I., Zapata, J. L., Dodelin, B., Alcázar, E., Barrios, V., Verdugo, A., Audisio, P., Micó, E., Otero, J. C., Bahillo, P., Viñolas, A., Valladares, L., Méndez, M., El Antry, S. & Galante, E. (2018) The Conservation Status and Distribution of Mediterranean Saproxylic Beetles. Malaga, Spain: IUCN. xii + 58. <https://doi.org/10.2305/iucn.ch.2018.ra.3.en>
- Gomy, Y., Labrique, H., Chavanon, G., Janati Idrissi, A. & François, A. (2011) Contribution à la connaissance des Histeridae du Maroc (Coleoptera). *Publications du Musée des Confluences*, 2(1), 23-74. <https://doi.org/10.3406/mhnly.2011.1526>
- Gosselin, M. & Pailland, Y. (2017) Mieux intégrer la biodiversité dans la gestion forestière Guide pratique (France métropolitaine). *Cemagref*, Editions QUAE, 100.
- Gultekin, L. & Fremuth, J. (2013) Lixinae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 8. Leiden, Brill, 437-471.
- Habbaz, H., Maatouf, N., Lumaret, J. P. & Rohi, L. (2025) The beetles (Coleoptera) associated with the Maamora cork oak forest in the north-west of Morocco. *Journal of Insect Biodiversity and Systematics*, 11(1), 39-84. <https://doi.org/10.61186/jibs.11.1.39>
- Hansen, M. (2004) Hydrophilidae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 2. Apollo Books, Stenstrup, 44-68.
- Hava, J. (2007a) Dermestidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 299-302.
- Hava, J. (2007b) Dermestidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 306-319.
- Hlavac, P. & Maughan, N. (2013) Cossoninae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 8. Leiden, Brill, 223-224.
- Horak, J. (2008) Mordellidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 87-105.
- Hutcheson, J. (1990) Characterization of terrestrial insect communities using quantified, Malaise-trapped Coleoptera. *Ecological Entomology*, 15(2), 143-151. <https://doi.org/10.1111/j.1365-2311.1990.tb00795.x>

- Iorio, É., Dusoulrier, F., Soldati, F., Noël, F., Guilloton, J.A., Doucet, G., Ponel, P., Dupont, P., Krieg-Jacquier, R., Chemin, S. & Tillier, P. (2022) Les Arthropodes terrestres dans les études d'impact: limites actuelles et propositions pour une meilleure prise en compte des enjeux de conservation. *Naturae*, 4, 43-99. <https://doi.org/10.5852/naturae2022a4>
- Îto, N. (2003) Carabidae: subtribe Anisodactylina. In: Löbl, I. & Smetana, A. (eds), Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 360-362.
- Iwan, D. & Löbl, I. (2008) Tenebrionidae: Tribe Melanimini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 257-258. https://doi.org/10.1163/9789004260900_012
- Jäch, M.A. (2004) Hydraenidae. In: Löbl, I. & Smetana, A. (eds), Catalogue of Palaearctic Coleoptera, 2. Apollo Books, Stenstrup, 102-122.
- Jactel, H., Imler, J. L., Lambrechts, L., Failloux, A. B., Lebrandon, J. D., Le Maho, Y., Duplessy, J. C., Cossart P. & Grandcolas, P. (2020) Insect decline: immediate action is needed. *Comptes Rendus. Biologie*, 343(3), 267-293. <https://doi.org/10.5802/crbio.37>
- Jaeger, B. & Kataev, B. M. (2003) Carabidae: subtribe, Stenolophina. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 397-406.
- Jeannel, R. (1941) Coléoptères Carabiques (première partie). *Faune de France*, 39, 571.
- Jelinek, J. & Audisio, P. (2007) Nitidulidae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 459-491.
- Jendek, E. (2006) Buprestidae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 388-421.
- Johnson, C. (2007) Latridiinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 635-641.
- Kabak, I. (2003) Carabidae: Tribe Lebiini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 408-438.
- Kang, S. H. & Lee, J. E. (2017) Morphological and Ecological study of Genus *Cryptocephalus* (Coleoptera: Chrysomelidae: Cryptocephalinae) from Korea. *국립공원연구지*, 8(3), 138-143.
- Karmaoui, A., Chavanon, G. & Moumane, A. (2023) Coleopteran Species in Arid Environments: Ecological Importance and Damages in the Pre-Sahara of Morocco. *Climate Change and the Economic Importance and Damages of Insects*, 119-145. <https://doi.org/10.4018/978-1-6684-4824-3.ch006>
- Kataev, B. M., Wrase, D. W. & Ito, N. (2003) Carabidae: Subtribe Harpalina. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 367-407.
- Kippenberg, H. (2010) Chrysomelinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 6. Apollo Books, Stenstrup, 390-443.
- Knizek, M. (2011) Scolytinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 7. Apollo Books, Stenstrup, 204-250.
- Kodada, J. & Jäch, M. A. (2006) Dryopidae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 440-443.
- Kolibac, J. (2007) Trogossitidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 364-366.
- Kovar, I. (2007) Coccinellidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 568-631.
- Kubán, V. (2006) Buprestinae without Anthaxini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 369-387.
- Leblanc, P., Levey, B. & Horak, J. (2008) Scaptiidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 458-465.
- Lefèvre, E. A. E., Fairmaire, L., Marseul, S. A. D. & Sénac, H. (1885) Exploration scientifique de la Tunisie publiée sous les auspices du ministère de l'Instruction publique. Sciences Naturelles, Zoologie - Coléoptères. Liste des Coléoptères recueillis en Tunisie en 1883 par M.A. Landourneux. Imprimerie Nationale, Paris. <https://doi.org/10.5962/bhl.title.53766>
- Lillig, M., Borg Barthet, H. & Mifsud, D. (2012) An identification and informative guide to the Tenebrionidae of Malta (Coleoptera). *Bulletin of the Entomological Society of Malta*, 5, 121-160. <https://www.um.edu.mt/library/oar/handle/123456789/538>

- Löbl, I., Krell, F. T., Kral, D. & Ziani, S. (2006a) Scarabaeidae: Tribe Onthophagini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 159–175.
- Löbl, I., Krell, F. T. & Kral, D. (2006b) Scarabaeidae: Tribe Scarabaeini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 176–177.
- Löbl, I., Merkl, O., Ando, K., Bouchard, P., Lillig, M., Masomuto, K. & Schawaller, W. (2008) Tenebrionidae without Lagriini, Asidini, Blaptini, Melanimini, Opatrini, Pedinini, Platyscelidini & Helopini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 105–352.
- Löbl, I., Nekolajev, G. V. & Kral, D. (2006c) Geotrupidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 84–92.
- Löbl, I. & Smetana, A. (2003) Archostemata-Myxophaga-Adephaga. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 1, 819. <https://doi.org/10.1086/382416>
- Löbl, I. & Smetana, A. (2004) Hydrophiloidea-Histeroidea-Staphylinoidea. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 2, 942. <https://doi.org/10.1086/497217>
- Löbl, I. & Smetana, A. (2006) Scarabaeoidea, Scirtoidea, Dascilloidea, Buprestoidea, Byrrhoidea. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 3, 690. [https://doi.org/10.1649/0010-065x\(2007\)61\[427:copcv\]2.0.co;2](https://doi.org/10.1649/0010-065x(2007)61[427:copcv]2.0.co;2)
- Löbl, I. & Smetana, A. (2007) Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 4, 935. https://doi.org/10.1163/9789004260894_004
- Löbl, I. & Smetana, A. (2008) Tenebrionoidea. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 4, 670. https://doi.org/10.1163/9789004260900_011
- Löbl, I. & Smetana, A. (2010) Chrysomeloidea. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 6, 924.
- Löbl, I. & Smetana, A. (2011) Curculionoidea I. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 7, 373. https://doi.org/10.1163/9789004260931_012
- Löbl, I. & Smetana, A. (2013) Curculionoidea II. Catalogue of Palaearctic Coleoptera. Stenstrup, Denmark, Apollo Books, 8, 700. https://doi.org/10.1163/9789004259164_010
- Lopatin, I., Smetana, A. & Schöller, M. (2010) Cryptocephalinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 6. Apollo Books, Stenstrup, 564–619.
- Lumbierres, B., Madeira, F. & Pons, X. (2018) Prey acceptability and preference of *Oenopia conglobata* (Coleoptera: Coccinellidae), a candidate for biological control in urban green areas. *Insects*, 9(1), 7. <https://doi.org/10.3390/insects9010007>
- Machouri, N. (2009) Les subéraies Marocaines face aux changements climatiques and actions anthropiques. Actes du colloque international de Niamey (Niger), pp. 297–312.
- Marggi, W. A., Huber, C., Müller-Motzfeld, G. & Hartmann, M. (2003) Carabidae, Tribe Bembidiini; subtribe Bembidiina. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 241–273.
- Mayor, A. (2007a) Dasytidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 388–415.
- Mayor, A. (2007b) Melyridae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 386–388.
- Mazur, S. (2004) Histeridae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 2. Apollo Books, Stenstrup, 68–102.
- Médail, F. & Quézel P. (1997) Hot-spots analysis for conservation of plant biodiversity in the Mediterranean basin. *Annals of the Missouri Botanical Garden*, 84, 112–127. <https://doi.org/10.2307/2399957>
- Micó, E. (2018) Saproxyllic insects in tree hollows. *Saproxyllic Insects: Diversity, Ecology and Conservation*, 1, 693–727. https://doi.org/10.1007/978-3-319-75937-1_21
- Nageleisen, L. M. & Bouget, C. (2009) L'étude des insectes en forêt : méthodes et techniques, éléments essentiels pour une standardisation. Synthèse des réflexions menées par le groupe de travail «Inventaires Entomologiques en Forêt». ONF, 144 p.
- Nikodym, M. & Bezdek, A. (2006) Glaphyridae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 97–103.
- Novak, V. & Pettersson, R. (2008) Tenebrionidae: Alleculinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 319–338.

- Numa, C., Tonelli, M., Lobo, J. M., Verdú, J. R., Lumarand, J. P., Sánchez-Piñero, F., Ruiz, J. L., Dellacasa, M., Ziani, S., Arriaga, A., Cabrero, F., Labidi, I., Barrios, V., Şenyüz, Y. & Anlaş, S. (2020) The conservation Status and Distribution of Mediterranean Dung Beetles. Gland, Switzerland and Málaga, Spain: IUCN.xii + 55. <https://doi.org/10.2305/iucn.ch.2020.ra.1.en>
- Pan, Z. & Bologna, M. A. (2014) Taxonomy, Bionomics and Faunistics of the Nominated Subgenus of *Mylabris* Fabricius, 1775, with the description of five new species (Coleoptera: Meloidae: Mylabrini). *Zootaxa*, 3806(1), 1-78. <https://doi.org/10.11646/zootaxa.3806.1.1>
- Parisi, F., Lombardi, F., Marziliano, P. A., Russo, D., De Cristofaro, A., Marchetti, M. & Tognetti, R. (2020) Diversity of saproxylic beetle communities in chestnut agroforestry systems. *Iforest-Biogeosciences and Forestry*, 13(5), 456. <https://doi.org/10.3832/for3478-013>
- Paulian, R. (1940) Coléoptères Scarabéides. *Faune de France*, 38, 239.
- Pellandier, J. (2013) Tribe Brachyderini. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, Vol. 8. Leiden, Brill, pp. 255-263.
- Peñuelas, J. & Sardans, J. (2021) Global change and forest disturbances in the Mediterranean basin: Breakthroughs, knowledge gaps, and recommendations. *Forests*, 12(5), 603. <https://doi.org/10.3390/f12050603>
- Puchkov, A. V. & Matalin, A. V. (2003) Carabidae, subfamily Cicindelinae, tribe Cicindelini. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 1. Apollo Books, Stenstrup, 99-115.
- Ramli, N.H., Rahim, N. A. A., Din, A. M. M. & Yaakop, S. (2019) A Preliminary Study on the Diversity and Abundance of Onthophagus Species (Coleoptera: Scarabaeidae) in an Oil Palm Plantation, Peninsular Malaysia in Relation to Carbon Dioxide and Soil Organic Matter. *Pertanika Journal of Tropical Agricultural Science*, 42(1), 405.
- Roger, J. L., Jambon, O. & Bouger, G. (2013) Clé de détermination des Carabides Paysages agricoles du Nord Ouest de la France. *INRA SAD*, 265.
- Ruzzier, E., Ghahari, H. & Horák, J. (2017) A checklist of the Iranian Mordellidae (Coleoptera: Tenebrionoidea). *Zootaxa*, 4320(1), 146-158. <https://doi.org/10.11646/zootaxa.4320.1.8>
- Sacco-Martret de Prévile, A., Staudacher, K., Traugott, M., Bohan, D. A., Plantegenest, M. & Canard, E. (2024) Prey Switching and Natural Pest Control Potential of Carabid Communities over the Winter Wheat Cropping Season. *Insects*, 15(8), 2075-4450. <https://doi.org/10.3390/insects15080610>
- Sallé, A., Cours, J., Le Souchu, E., Lopez-Vaamonde, C., Pincebourde, S. & Bouget C. (2021) Climate Change Alters Temperate Forest Canopies and Indirectly Reshapes Arthropod Communities. *Frontiers in Forests and Global Change*, 4, 710854. <https://doi.org/10.3389/ffgc.2021.710854>
- Sama, G. (1987) Note préliminaire pour une faune des Longicornes de l'Afrique-du-Nord (Coleoptera, Cerambycidae). *Biocosme Mesogéen*, 4(1), 43-64.
- Sama, G. & Löbl, I. (2010a) Cerambycinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 6. Apollo Books, Stenstrup, 143-206.
- Sama, G. & Löbl, I. (2010b) Lepturinae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 6. Apollo Books, Stenstrup, 114-117.
- Samih, A., Trócoli, S., Rohi, L., Fougrach, H., Hsaine, M. & Maatouf, N. (2024) Comparative study of the diversity and structure of plant-pollinator interactions in forest and agricultural landscapes in Northwestern Morocco. *European Journal of Entomology*, 121, 400-412. <https://doi.org/10.14411/eje.2024.044>
- Sauvage, C. (1961) Recherches géobotaniques sur les subéraies Marocaines. *Travaux de l'Institut Scientifique Chérifien, série Botanique*, 21, 1-452.
- Secchi, F. (2002) Histeridae de France (Coleoptera, Histeridae). Supplément au Bulletin de la Société Linnéenne de Bordeaux, 19(4), 433-472.
- Seibold, S. & Thorn, S. (2018) The importance of dead-wood amount for saproxylic insects and how it interacts with dead-wood diversity and other habitat factors. *Saproxylic Insects: Diversity, Ecology and Conservation*, 607-637. https://doi.org/10.1007/978-3-319-75937-1_18
- Sforzi, A. (2011) Brentidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 7. Apollo Books, Stenstrup, 142-148.
- Sheikh, A. H., Thomas, M., Bhandari, R. & Meshram, H. (2016) Malaise trap and insect sampling: Mini Review. *Bio Bulletin*, 2(2), 35-40.
- Šlipinski, S. A. & Schuh, R. (2008) Zopheridae. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 548-552.

- Smetana, A. (2004) Staphylinidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 2. Apollo Books, Stenstrup, 237-699.
- Smetana, A. (2006) Melolonthidae: Tribe Hopliini. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 184-190.
- Schmidt, M. H., Thewes, U., Thies, C. & Tscharnkte, T. (2004) Aphid suppression by natural enemies in mulched cereals. *Entomologia Experimentalis et Applicata*, 113(2), 87-93. <https://doi.org/10.1111/j.0013-8703.2004.00205.x>
- Švec, Z. (2007) Phalacridae. In: I. Löbl & A. Smetana (eds.) Catalogue of Palaearctic Coleoptera, 4. Appolo Books, Stenstrup, 507-513.
- Švihla, V. (2008) Oedemeridae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 5. Apollo Books, Stenstrup, 353-369.
- Telnov, D. M. I. T. R. Y. (2010) Ant-like flower beetles (Coleoptera: Anthicidae) of the UK, Ireland and Channel Isles. *British Journal of Entomology and Natural History*, 23(2), 99.
- Thotagamuwa, A., Noriega, J. A., Webb, S., Weston, P., Doube, B. M., Caron, V., Gleeson, P., Forgie, S. A., Favila, M. E., Burgess, G. & Gurr, G. M. (2023) Rearing dung beetles (Coleoptera: Scarabaeidae): identifying knowledge gaps and future challenges. *Entomologia Generalis*, 43(4), 751-769. <https://doi.org/10.1127/entomologia/2023/1716>
- Tronquet, M. (2021) Catalogue des Coléoptères de France, 917.
- Trócoli Gracia, S. (2018) Actualización del catálogo de Longicornios de Marruecos (Parte I: Prioninae, Aseminae, Saphaninae). *Revue de l'Association Roussillonnaise d'Entomologie*, Tome XXVII (3), 130-140.
- Trócoli Gracia, S. (2019) Actualización del catálogo de Longicornios de Marruecos (Parte II: Cerambycidae: Lepturinae, Vesperidae). *Revue de l'Association Roussillonnaise d'Entomologie*, Tome XXVIII (2), 72-84.
- Trócoli Gracia, S. (2020) Actualización del catálogo de Longicornios de Marruecos Actualisation du catalogue des Longicornes du Maroc (Parte IV: Cerambycidae: Lamiinae). *Revue de l'Association Roussillonnaise d'Entomologie*, Tome XXIX (1), 26-65.
- Trócoli, S. (2023) Nuevas aportaciones al Catálogo de Longicornios de Marruecos. *Revue de l'Association Roussillonnaise d'Entomologie*, 32(3), 40. <http://hdl.handle.net/11703/131001>
- Tsiakiris, R., Stara, K., Kazoglou, Y., Kakouros, P., Bousbouras, D., Dimalexis, A., Dimopoulos, P., Fotiadis, G., Gianniris, I., Kokkoris, I. P. & Mantzanas, K. (2024) Agroforestry and the Climate Crisis: Prioritizing Biodiversity Restoration for Resilient and Productive Mediterranean Landscapes. *Forests*, 15(9), 1648. <https://doi.org/10.3390/f15091648>
- Tsikas, A. & Karanikola, P. (2022) To conserve or to control? Endangered saproxylic beetles considered as forest pests. *Forests*, 13(11), 1929. <https://doi.org/10.3390/f13111929>
- Ulyshen, M. D., Hanula, J. L. & Horn, S. (2005) Using Malaise traps to sample ground beetles (Coleoptera: Carabidae). *The Canadian Entomologist*, 137(2), 251-256. <https://doi.org/10.4039/n04-035>
- Velazquez de Castro, A. J. (2013) Tribe Sitonini. In: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, 8. Leiden, Brill, 386-392.
- Verdu, J. R. & Galante, E. (2006) Libro Rojo de Los Invertebrados de España. Dirección General de Conservación de la Naturaleza, Madrid.
- Verdu, J. R. & Galante, E. (2009) Atlas de los Invertebrados Amenazados de España (Especies en Peligro Crítico y en Peligro). Dirección General para la Biodiversidad, Ministerio de Medio Ambiente, Madrid, 340.
- Villemant, C. & Fraval, A. (1993) The insect fauna of the cork-oak tree in the Mamora forest (Morocco). *Ecologia Mediterranea (France)*, 19(3), 89-98. <https://doi.org/10.3406/ecmed.1993.1725>
- Volkovitsh, M. G. (2006) Buprestidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 3. Apollo Books, Stenstrup, 325-342.
- Wegrzynowicz, P., (2007a) Erotylidae. In: Löbl, I. & Smetana, A. (eds): Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 89-209.
- Wegrzynowicz, P. (2007b) Laemophloeidae. In: Löbl, I. & Smetana, A. (eds.) Catalogue of Palaearctic Coleoptera, 4. Apollo Books, Stenstrup, 503-507.
- Xofis, P., Feloni, E., Emmanouloudis, D., Chatzigiovanakis, S., Kravari, K., Samourkasidou, E., Kefalas, G. & Nastos, P. (2024) Long-Term Monitoring of Trends in Xerothermality and Vegetation Condition of a Northeast Mediterranean Island Using Meteorological and Remote Sensing Data. *Land*, 13(12), 2129. <https://doi.org/10.3390/land13122129>
- Zerga, B., Warkineh, B., Teketay, D. & Woldetsadik, M. (2021) The sustainability of reforesting landscapes with exotic species: a case study of eucalypts in Ethiopia. *Sustainable Earth*, 4(1), 5. <https://doi.org/10.1186/s42055-021-00044-7>

Zumr, V., Nakládal, O., Gallo, J. & Remeš, J. (2024) Deadwood position matters: Diversity and biomass of saproxylic beetles in a temperate beech forest. *Forest Ecosystems*, 11, 100174. <https://doi.org/10.1016/j.fecs.2024.100174>

Citation: Amine, S., Noureddin, M., Sergi, T., Hamza, H. & Latifa, R. (2025) Checklist of beetles (Coleoptera) from the cork oak forest of Larache (Morocco). *J. Entomol. Soc. Iran* 45 (2), 181–225.

DOI: <https://doi.org/10.22034/jesi.45.2.3>

URL: https://jesi.areeo.ac.ir/article_133227.html



فهرست سفت‌بالپوشان جنگل بلوط‌پنبه لاراش، مراکش

امین سمیح^۱ , نورالدین معطوف^۲ , سرجی ترکولی^۳ , حمزه حجاز^۴  و لطیفه روحی^۱ 

۱- آزمایشگاه بوم‌شناسی و محیط‌زیست، دانشکده علوم بن مسیک، دانشگاه حسن دوم، کازابلانکا، مراکش

۲- مرکز نوآوری، پژوهش و آموزش، سازمان ملی آب و جنگل‌ها، رباط، مراکش

۳- موزه علوم طبیعی بارسلونا، آزمایشگاه طبیعت، کلکسیون بندپایان، بارسلونا، اسپانیا

چکیده: جنگل بلوط‌پنبه لاراش، که بخش حیاتی از جنگل‌های بلوط‌پنبه سواحل اقیانوس اطلس در مراکش است، نقش حیاتی در حمایت از اکوسیستم‌های محلی و جوامع انسانی ایفا می‌کند. با این حال، این جنگل به‌طور فزاینده‌ای در معرض چالش‌هایی چون خطر کشاورزی فشرده و دیگر دست‌اندازی‌های بشر قرار دارد که منجر به نابودی زیستگاه‌های حشرات از جمله جوامع سوسک‌ها شده است. هدف از این مطالعه، تهیه فهرستی جامع از سخت‌بالپوشان در جنگل بلوط‌پنبه لاراش در طی دو سال متوالی (۲۰۲۱ و ۲۰۲۲) بود که از طریق بررسی سوسک‌های مرتبط با چشم‌اندازهای گیاهی مختلف با استفاده از تکنیک‌های مختلف تله‌گذاری (تله‌های گودال و تله‌های کاسه‌ای) و بررسی‌های مستقیم انجام شد. نتایج نشان از حضور ۱۸۰ گونه سوسک متعلق به ۳۷ خانواده داشت که شامل ۹ گونه بومی، ۳۳ گونه که به تازگی در مراکش ثبت شده‌اند، ۱۳ گونه سابروکسیلیک و ۱۰ گونه فهرست‌شده در فهرست سرخ شامل هفت گونه خشک‌دارزی (سپروکسیلیک) و سه گونه سرگین خوار (کوپروفاگوس) هستند. این تنوع، اهمیت اکولوژیک جنگل بلوط‌پنبه لاراش را نشان می‌دهد و نیاز فوری به اعمال روش‌های حفاظتی و تقویت زیستگاه‌های مناسب برای حفاظت از این جوامع حشرات را برجسته می‌کند.

تاریخچه مقاله

دریافت: ۱۴۰۳/۰۹/۱۳

پذیرش: ۱۴۰۳/۱۱/۲۷

انتشار: ۱۴۰۴/۰۱/۲۴

دبیر تخصصی: مار فر سوای

نویسنده مسئول: امین سمیح

ایمیل: aminesamih96@gmail.com

DOI: <https://doi.org/10.22034/jesi.45.2.3>

کلمات کلیدی: حفاظت، نمونه برداری، تنوع، آفریقا، حاشیه اقیانوس اطلس